


## 2-Intertidal and Nearshore Benthos

### Introduction

A major objective of the Biological Monitoring Program (BMP) is the detection of changes in intertidal benthic assemblages as a consequence of fill operations. Inherent limitations in the power of sampling programs to detect less than dramatic changes in fishery resources underscore the general reliance on quantitatively sampling benthos in order to infer impacts on fishery resources. Impacts to beach infauna are expected to include reduced abundance and altered community structure. While there are no standard sampling programs for collecting this type of information, guidelines found in Cochran (1963), Morrissey et al. (1992), and Nelson (1993) have been followed. Likewise, recommendations concerning statistical design found in Saila et al. (1976), Cohen (1988), and Underwood (1992) were applied to the study. Emphasis is placed on detecting both short-term and long-term impacts to abundance, biomass, and assemblage structure along a gradient from intertidal to subtidal depths.

Examination of nourishment impacts to New Jersey beaches is complicated by the relative paucity of quantitative studies of beach infauna along the mid-Atlantic coast. In a review of infaunal zonation on high-energy beaches McLachlan and Jaramillo (1995) reported no studies between North Carolina and southern New England. Fortunately, McDermott (1983) has described the distribution of southern New Jersey beach infauna as part of a study of the surf zone food web. New Jersey beach infauna were dominated by the polychaete *Scolecopsis squamata*, the wedge clam *Donax variabilis*, the mole crab *Emerita talpoida*, the ribbon worm (Rhynchocoela) *Micura leidy*, and a variety of haustoriid amphipods. McDermott has previously reported the distribution of both *S. squamata* (McDermott, 1979) and the nephtyid polychaete *Nephtys bucera* (McDermott, 1987) from the same site. Croker (1970) provides a faunal list of intertidal macrofauna from Long Island that is very similar to that of McDermott (1983). The benthic assemblage described in these reports is essentially identical to these and others reported for U.S Atlantic coast beaches (McLachlan and Jaramillo, 1995).

In addition to the paucity of reports concerning the quantitative distribution of beach infauna, there are virtually no studies of beach nourishment impacts to intertidal infauna north of the Carolinas. Both Nelson (1985 and 1993) and Hackney et al. (1996) have extensively reviewed the literature and neither report studies for this region. The closest study is that of Jaramillo et al. (1987) who followed the recovery of a New England sandy beach after intense erosion. In general, both Nelson (1985 and 1993) and Hackney et al. (1996) categorized nourishment impacts to beach infauna as short-term with recovery times ranging from 2 to 7 months. Subsequent studies by Jutte et al. (1999a and 1999b) in South Carolina arrived at a similar estimate (3 to 6 months). The longest recovery times reported for nourished beaches occur when there is a poor match between fill materials and the original substrate. For instance, Reilly and Bellis (1983) reported that recovery took more than a year for some species on a North Carolina beach where substantial amounts of silts and clays were present in the fill.



**Figure 2-1. Intertidal and Nearshore benthos sampling locations.**

**Table 2-1. Summary Biological Results for Intertidal and Nearshore Infauna**

Data	Area	Total			South			Middle			North		
	Depth	MLW	MLW-1m	Nearshore									
Total Taxa		80	80	225	57	54	173	42	48	137	41	45	158
Total Animals		81691	50827	298999	25633	16360	125827	38963	16581	35669	17095	17886	137503
Total Biomass (g)		779.68	502.87	32906.64	380.61	165.78	8554.47	236.32	184.55	17030.50	162.76	152.54	7321.67
Average Taxa/Sample*		2.6	2.5	14.3	2.7	2.4	15.9	2.6	2.5	13.0	2.5	2.7	14.2
Average Abundance/m <sup>2</sup>		11503	7272	632	10552	7097	721	16583	7262	250	7373	7458	883
Average Biomass (g/m <sup>2</sup> )		150.60	97.14	99.72	220.56	96.07	77.77	136.94	106.94	154.82	94.31	88.40	66.56
Annelid		107.69	75.58	1.44	150.40	75.22	1.40	105.71	81.58	1.02	66.95	69.93	1.88
Crustacea		21.36	19.10	9.05	29.39	15.79	9.03	11.78	24.47	14.57	22.91	17.05	3.54
Mollusc		20.73	0.63	88.97	39.87	0.41	67.16	18.12	0.40	138.96	4.19	1.07	60.79
Echinoderm		-----	4.21	---	---	4.21	---	---	---	---	---	---	0.00
Miscellaneous		0.80	0.42	0.27	0.87	0.42	0.17	1.26	0.48	0.27	0.26	0.35	0.35
% Annelid		72.12	77.90	10.12	68.19	78.30	5.53	77.19	76.29	5.71	70.98	79.11	19.11
% Crustacean		15.41	19.54	24.33	13.32	16.44	26.56	8.61	22.88	23.87	24.29	19.29	22.56
% Mollusc		11.92	0.67	63.50	18.07	0.43	66.97	13.23	0.38	68.62	4.44	1.21	54.91
% Echinoderm		-----	4.38	0.00	---	4.38	0.00	---	---	0.01	---	---	0.00
% Miscellaneous		0.53	0.43	2.07	0.39	0.44	0.94	0.92	0.45	1.80	0.28	0.40	3.49

\*Intertidal sample = 44cm<sup>2</sup>; Nearshore sample = 0.1m<sup>2</sup>

**Appendix Table 2-1. Mean Low Water Station Locations**

<b>Area</b>	<b>Station No.</b>	<b>Depth</b>	<b>Latitude</b>	<b>Longitude</b>
South	1	MLW	40° 08.006	74° 01.665
South	2	MLW	40° 08.139	74° 01.621
South	3	MLW	40° 08.588	74° 01.465
South	4	MLW	40° 08.743	74° 01.402
South	5	MLW	40° 08.960	74° 01.343
South	6	MLW	40° 09.064	74° 01.376
South	7	MLW	40° 09.263	74° 01.273
South	8	MLW	40° 09.397	74° 01.251
South	9	MLW	40° 09.579	74° 01.085
South	10	MLW	40° 09.701	74° 01.115
Middle	21	MLW	40° 11.377	74° 00.554
Middle	22	MLW	40° 11.478	74° 00.527
Middle	23	MLW	40° 11.568	74° 00.509
Middle	24	MLW	40° 11.798	74° 00.440
Middle	25	MLW	40° 12.004	74° 00.426
Middle	26	MLW	40° 12.198	74° 00.361
Middle	27	MLW	40° 12.454	74° 00.285
Middle	28	MLW	40° 12.545	74° 00.216
Middle	29	MLW	40° 12.803	74° 00.097
Middle	30	MLW	40° 12.936	74° 00.056
North	41	MLW	40° 14.091	73° 59.724
North	42	MLW	40° 14.147	73° 59.672
North	43	MLW	40° 14.588	73° 59.610
North	44	MLW	40° 14.645	73° 59.576
North	45	MLW	40° 14.982	73° 59.480
North	46	MLW	40° 15.019	73° 59.470
North	47	MLW	40° 15.080	73° 59.408
North	48	MLW	40° 15.100	73° 59.351
North	49	MLW	40° 15.296	73° 59.414
North	50	MLW	40° 15.340	73° 59.353

**Appendix Table 2-2. Abundance of Intertidal Taxa (Numbers of Animals/m<sup>2</sup>)**

Taxon (LPIL)	Area	South	South											
	Depth	MLW	MLW											
	Date	May-94	Sept.- 94	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
Acarina (LPIL)														
Actiniaria (LPIL)														
<i>Ampelisca abdita</i>				2486		226								
Ampharetidae (LPIL)														
Amphipoda (LPIL)										226				
<i>Amphiporeia gigantea</i>														
<i>Ampithoe valida</i>						226								
<i>Ancistrosyllis hartmanae</i>														
Archannelida Family A				339										
Ascidacea (LPIL)														
Bivalvia (LPIL)									226	226				
Bodotriidae (LPIL)						226								
Branchiopoda (LPIL)														
Calyptraeidae (LPIL)														
<i>Cancer irroratus</i>														
Capitellidae (LPIL)														
<i>Chiridotea tuftsi</i>														
Cirratulidae (LPIL)				1695										
Cladocera (LPIL)														
<i>Corophium tuberculatum</i>														
<i>Crangon septemspinosa</i>														
<i>Crepidula</i> (LPIL)									904					
<i>Crepidula fornicata</i>														
<i>Crepidula plana</i>								226						
Decapoda (LPIL)			226											
Decapoda Reptantia (LPIL)														
<i>Dipolydora socialis</i>														
<i>Donax variabilis</i>										301			339	
Echinoidea (LPIL)														
<i>Emerita talpoida</i>		226	291	678	226	1582	226	452		881	678	678	904	
<i>Eteone longa</i>										226				

**Appendix Table 2-2. Abundance of Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxon (LPIL)	Area	South	South											
	Depth	MLW	MLW											
	Date	May-94	Sept.- 94	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Eurythoe complanata</i>														
Gammaridae (LPIL)						226								
<i>Gammarus annulatus</i>			226			452								
Gastropoda (LPIL)										226				
<i>Gemma gemma</i>														
Goneplacidae (LPIL)														
<i>Goniada teres</i>													226	
Haustoriidae (LPIL)														
<i>Haustorius canadensis</i>								226			226			226
<i>Haustorius sp. A</i>			226											
<i>Haustorius sp. C</i>														
Hesionidae (LPIL)			226											
Hirudinea (LPIL)			1582											
<i>Jassa falcata</i>									226			226		226
<i>Levinsenia gracilis</i>							301							
<i>Litocorsa antennata</i>							226							
Lumbrineridae (LPIL)								226						
Mactridae (LPIL)									452					
<i>Magelona annulata</i>														
<i>Magelona papillicornis</i>														
<i>Mediomastus ambiseta</i>		226			565		226							
Melitidae (LPIL)														
<i>Microphthalmus (LPIL)</i>					226									
<i>Microphthalmus aberrans</i>			226		16046									
<i>Microphthalmus hartmanae</i>														
<i>Microphthalmus sp. G</i>					603	565				339			396	
<i>Microphthalmus listensis</i>														
<i>Mysella planulata</i>														
Mysidacea (LPIL)														
Mytilidae (LPIL)								226						
<i>Mytilus edulis</i>		517	1098	1281	19210	226	452	1456	5368	2769		7860		678







**Appendix Table 2-2. Abundance of Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxon (LPIL)	Area	South	South											
	Depth	MLW-1	MLW-1											
	Date	May-94	Sept.- 94	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Eurythoe complanata</i>														
Gammaridae (LPIL)														
<i>Gammarus annulatus</i>														226
Gastropoda (LPIL)														
<i>Gemma gemma</i>				226										
Goneplacidae (LPIL)														
<i>Goniada teres</i>														
Haustoriidae (LPIL)														
<i>Haustorius canadensis</i>														301
<i>Haustorius sp. A</i>			377											
<i>Haustorius sp. C</i>		226												
Hesionidae (LPIL)										377				
Hirudinea (LPIL)														
<i>Jassa falcata</i>									452					
<i>Levinsenia gracilis</i>														
<i>Litocorsa antennata</i>														
Lumbrineridae (LPIL)														
Mactridae (LPIL)														
<i>Magelona annulata</i>							226							
<i>Magelona papillicornis</i>							452							
<i>Mediomastus ambiseta</i>				678			904							
Melitidae (LPIL)														226
<i>Microphthalmus (LPIL)</i>		904	226	226		452				1130				
<i>Microphthalmus aberrans</i>			678		1921									
<i>Microphthalmus hartmanae</i>														
<i>Microphthalmus sp. G</i>						1582	1130			1017				
<i>Microphthalmus listensis</i>						226								
<i>Mysella planulata</i>														
Mysidacea (LPIL)							226							
Mytilidae (LPIL)														
<i>Mytilus edulis</i>		565	362	226	226	226	226	1040	452	961		1883		283







**Appendix Table 2-2. Abundance of Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxon (LPIL)	Area	Middle	Middle											
	Depth	MLW	MLW											
	Date	May-94	Sept.- 94	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Eurythoe complanata</i>														
Gammaridae (LPIL)														
<i>Gammarus annulatus</i>														
Gastropoda (LPIL)										226				
<i>Gemma gemma</i>														
Goneplacidae (LPIL)														
<i>Goniada teres</i>														
Haustoriidae (LPIL)														
<i>Haustorius canadensis</i>								452					226	
<i>Haustorius sp. A</i>														
<i>Haustorius sp. C</i>		452			226						339			
Hesionidae (LPIL)														
Hirudinea (LPIL)			226											
<i>Jassa falcata</i>								226		226				
<i>Levinsenia gracilis</i>														
<i>Litocorsa antennata</i>														
Lumbrineridae (LPIL)														
Mactridae (LPIL)														
<i>Magelona annulata</i>														
<i>Magelona papillicornis</i>														
<i>Mediomastus ambiseta</i>							226							
Melitidae (LPIL)														
<i>Microphthalmus (LPIL)</i>		452	452	339	1507		452	7458			678	1582		
<i>Microphthalmus aberrans</i>			226		1281									
<i>Microphthalmus hartmanae</i>														
<i>Microphthalmus sp. G</i>						226	452	2260	1582	452				
<i>Microphthalmus listensis</i>														
<i>Mysella planulata</i>										226				
Mysidacea (LPIL)														
Mytilidae (LPIL)													226	
<i>Mytilus edulis</i>		414	3239	226		226	452	1065	735	1780		1306	226	452







**Appendix Table 2-2. Abundance of Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxon (LPIL)	Area	Middle	Middle											
	Depth	MLW-1	MLW-1											
	Date	May-94	Sept.- 94	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Eurythoe complanata</i>														
Gammaridae (LPIL)														
<i>Gammarus annulatus</i>														
Gastropoda (LPIL)													226	
<i>Gemma gemma</i>														
Goneplacidae (LPIL)										226				
<i>Goniada teres</i>														
Haustoriidae (LPIL)		226	226	226				226				226		
<i>Haustorius canadensis</i>								226						
<i>Haustorius sp. A</i>														
<i>Haustorius sp. C</i>			226		678						226			
Hesionidae (LPIL)														
Hirudinea (LPIL)														
<i>Jassa falcata</i>								226				226	226	
<i>Levinsenia gracilis</i>														
<i>Litocorsa antennata</i>														
Lumbrineridae (LPIL)														
Mactridae (LPIL)														
<i>Magelona annulata</i>														
<i>Magelona papillicornis</i>														
<i>Mediomastus ambiseta</i>					3239									
Melitidae (LPIL)														
<i>Microphthalmus (LPIL)</i>			339	452		678				7684				
<i>Microphthalmus aberrans</i>					1055									
<i>Microphthalmus hartmanae</i>														
<i>Microphthalmus sp. G</i>										452				
<i>Microphthalmus listensis</i>						1017								
<i>Mysella planulata</i>														
Mysidacea (LPIL)														
Mytilidae (LPIL)											226			
<i>Mytilus edulis</i>		1130	301	264	226	339	226	1384	791	1424	226	904		226







**Appendix Table 2-2. Abundance of Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxon (LPIL)	Area	North	North											
	Depth	MLW	MLW											
	Date	May-94	Sept.- 94	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Eurythoe complanata</i>														
Gammaridae (LPIL)														
<i>Gammarus annulatus</i>				226										
Gastropoda (LPIL)								226						
<i>Gemma gemma</i>														
Goneplacidae (LPIL)														
<i>Goniada teres</i>														
Haustoriidae (LPIL)											226			
<i>Haustorius canadensis</i>														
<i>Haustorius sp. A</i>														
<i>Haustorius sp. C</i>			452											
Hesionidae (LPIL)														
Hirudinea (LPIL)			4219											
<i>Jassa falcata</i>														
<i>Levinsenia gracilis</i>														
<i>Litocorsa antennata</i>														
Lumbrineridae (LPIL)														
Mactridae (LPIL)														
<i>Magelona annulata</i>														
<i>Magelona papillicornis</i>														
<i>Mediomastus ambiseta</i>				2712										
Melitidae (LPIL)														
<i>Microphthalmus (LPIL)</i>		226	1959	2034		791			1356		565	2185	791	
<i>Microphthalmus aberrans</i>		753	1808								904	1808		
<i>Microphthalmus hartmanae</i>														
<i>Microphthalmus sp. G</i>						678	10848		226	2712			1582	
<i>Microphthalmus listensis</i>														
<i>Mysella planulata</i>														
Mysidacea (LPIL)														
Mytilidae (LPIL)											226			
<i>Mytilus edulis</i>		716	497	509		603	565	778	527	1092		2283	226	1413







**Appendix Table 2-2. Abundance of Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxon (LPIL)	Area	North	North											
	Depth	MLW-1	MLW-1											
	Date	May-94	Sept.- 94	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Eurythoe complanata</i>						226								
Gammaridae (LPIL)														
<i>Gammarus annulatus</i>					226									
Gastropoda (LPIL)												226		
<i>Gemma gemma</i>														
Goneplacidae (LPIL)														
<i>Goniada teres</i>														
Haustoriidae (LPIL)			226						452		226			
<i>Haustorius canadensis</i>														
<i>Haustorius sp. A</i>														
<i>Haustorius sp. C</i>			226								226			
Hesionidae (LPIL)														
Hirudinea (LPIL)		1733												
<i>Jassa falcata</i>										226				
<i>Levinsenia gracilis</i>														
<i>Litocorsa antennata</i>														
Lumbrineridae (LPIL)														
Mactridae (LPIL)														
<i>Magelona annulata</i>														
<i>Magelona papillicornis</i>														
<i>Mediomastus ambiseta</i>														
Melitidae (LPIL)														
<i>Microphthalmus (LPIL)</i>			1582	1733		6272				3345	678			
<i>Microphthalmus aberrans</i>		9718									2260			
<i>Microphthalmus hartmanae</i>		1356	4068											
<i>Microphthalmus sp. G</i>						4972				6504				
<i>Microphthalmus listensis</i>						904								
<i>Mysella planulata</i>														
Mysidacea (LPIL)														
Mytilidae (LPIL)														
<i>Mytilus edulis</i>		1620	339	377		226	226	1243	377	1758		1893	226	2893





**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>)**

Taxa	Area	South	South	South								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Acanthohaustorius</i> (LPIL)				10	50		70					
<i>Acanthohaustorius intermedius</i>												
<i>Acanthohaustorius millsi</i>		48	119	55	251	236	477	235	611	337	384	318
Acarina (LPIL)												
Actiniaria (LPIL)				10	10		40		10	120		
<i>Albunea paretii</i>							10					
<i>Americamysis bigelowi</i>					10		70	10				
<i>Americhelidium americanum</i>		13	18	20			10	30	10	10		10
<i>Ampelisca</i> (LPIL)							10					
<i>Ampharete</i> (LPIL)			50	28								
<i>Ampharete americana</i>		30	20	50		22						
<i>Ampharete finmarchica</i>				50								
Ampharetidae (LPIL)		15	58	23		40	56	15	10	208	15	470
Amphipoda (LPIL)		10		10	10	10						
<i>Ampithoe</i> (LPIL)					20							
<i>Ampithoe longimana</i>												
<i>Ampithoe valida</i>					20							
<i>Anadara ovalis</i>							150					
<i>Ancinus depressus</i>												
Aoridae (LPIL)												
<i>Apoprionospio pygmaea</i>									10			
Archannelida Family A				10								
Arcidae (LPIL)												
<i>Aricidea</i> (LPIL)							20					
<i>Aricidea catherinae</i>												
<i>Asabellides oculata</i>						35	228	160		90	104	998
Ascidiacea (LPIL)												
<i>Astarte castanea</i>							338					
Asteroidea (LPIL)			50				140					
<i>Balanoglossus</i> (LPIL)												10
<i>Bathyporeia</i> (LPIL)					120							
<i>Bathyporeia parkeri</i>				10		41		10		10	20	

**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area Date	South May-95	South Sept.- 95	South May-96	South Sept.- 96	South May-97	South Sept.- 97	South May-98	South Sept.- 98	South May-99	South Sept.- 99	South May-00
<i>Bathyporeia quoddyensis</i>								10	30	30		38
Bivalvia (LPIL)							50	20	15			90
Bodotriidae (LPIL)												
<i>Callinectes</i> (LPIL)							10					
<i>Callinectes similis</i>							10					
<i>Calyptraea centralis</i>												
Calyptraeidae (LPIL)							80					
<i>Cancer irroratus</i>												
<i>Capitella capitata</i>				10						200		
Capitellidae (LPIL)										20	10	
<i>Carazziella hobsonae</i>				10								
<i>Caulleriella sp. J</i>										10		
<i>Chaetozone</i> (LPIL)			10									
<i>Chiridotea tuftsi</i>		178	10	59	133	194		75	102	71	22	24
Cirratulidae (LPIL)			10	10				10	10	40		20
<i>Corophium</i> (LPIL)								20				
<i>Corophium acutum</i>												
<i>Corophium insidiosum</i>												
<i>Corophium tuberculatum</i>								10			10	
<i>Crangon septemspinosa</i>			10		10			17	10			
<i>Crepidula</i> (LPIL)												
<i>Crepidula fornicata</i>								70				
<i>Crepidula plana</i>								128	10			
Cumacea (LPIL)												
Decapoda (LPIL)												
Decapoda Reptantia (LPIL)			10									
Diastylidae (LPIL)								10				
<i>Diastylis sculpta</i>									10			
<i>Diopatra cuprea</i>												
<i>Dipolydora commensalis</i>										10		
<i>Dipolydora socialis</i>		10		10								
<i>Dispia uncinata</i>		26	64	23	42			18	190	55	250	18







**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	South	South	South								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Paranaitis speciosa</i>						10						
Paraonidae (LPIL)												
<i>Paraonis fulgens</i>		17										
<i>Paraprionospio pinnata</i>				10								
<i>Pectinaria gouldii</i>							10					
Pelecypoda (LPIL)		10	10	54	30							
<i>Petricola pholadiformis</i>											20	50
<i>Pherusa affinis</i>							10					
Pholadidae (LPIL)		10										
<i>Photis macrocoxa</i>												
Phoxocephalidae (LPIL)												
<i>Phyllodoce</i> (LPIL)							10					
<i>Phyllodoce arenae</i>		20	10				29			20	10	28
Phyllodocidae (LPIL)												
<i>Pitar morrhuanus</i>										10	10	
Podocopida (LPIL)												
<i>Politilana polita</i>												
<i>Polydora cornuta</i>												
<i>Polygordius</i> (LPIL)			10		10			10	18		37	15
Polynoidae (LPIL)				10			60					
Portunidae (LPIL)							10					
<i>Protohaustorius</i> (LPIL)				25	10		30	130				
<i>Protohaustorius sp. B</i>		82	102	53	50	37	399	165	138			
<i>Protohaustorius wigleyi</i>										270	388	269
<i>Psammonyx nobilis</i>		107	207	1400	110	570	365	73	140	515	392	1215
<i>Pseudoleptocuma minor</i>					23	10		18				10
<i>Pseudunciola obliquua</i>			35						15			
Pyramidellidae (LPIL)							10					
<i>Rhepoxynius hudsoni</i>									10			
Rhynchozoela (LPIL)		28	15	20	24	20	25	14	10	28	14	20
<i>Sabellaria vulgaris</i>			10				251					
<i>Scolecopsis squamata</i>						15	25	25	305	90	10	



**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	Middle	Middle	Middle								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Acanthohaustorius</i> (LPIL)			15		33		35	20	15			
<i>Acanthohaustorius intermedius</i>		10										
<i>Acanthohaustorius millsi</i>		46	43	27	61	134	141	220	307	68	356	129
Acarina (LPIL)												
Actiniaria (LPIL)				320						33	10	
<i>Albunea paretii</i>												
<i>Americamysis bigelowi</i>			10		15	10	31		10			
<i>Americhelidium americanum</i>		10	14	10	10	10			15		10	
<i>Ampelisca</i> (LPIL)												
<i>Ampharete</i> (LPIL)			35	10								
<i>Ampharete americana</i>		15	53	10								
<i>Ampharete finmarchica</i>												
Ampharetidae (LPIL)		10	17	10	10	20	1155	40	10	32	23	10
Amphipoda (LPIL)					10							
<i>Ampithoe</i> (LPIL)			10									
<i>Ampithoe longimana</i>			10									
<i>Ampithoe valida</i>												
<i>Anadara ovalis</i>												
<i>Ancinus depressus</i>												10
Aoridae (LPIL)												
<i>Apoprionospio pygmaea</i>												
Archannelida Family A												
Arcidae (LPIL)												
<i>Aricidea</i> (LPIL)												
<i>Aricidea catherinae</i>												
<i>Asabellides oculata</i>						20	92	135		109	120	
Asciadiacea (LPIL)											30	
<i>Astarte castanea</i>							145	20				
Asteroidea (LPIL)							10		15			
<i>Balanoglossus</i> (LPIL)												
<i>Bathyporeia</i> (LPIL)												
<i>Bathyporeia parkeri</i>		10				25		10				

**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	Middle	Middle	Middle								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Bathyporeia quoddyensis</i>								10				
Bivalvia (LPIL)						10	50	10	10	20		10
Bodotriidae (LPIL)			10									
<i>Callinectes</i> (LPIL)												
<i>Callinectes similis</i>												
<i>Calyptraea centralis</i>												
Calyptraeidae (LPIL)				20			40					
<i>Cancer irroratus</i>												
<i>Capitella capitata</i>												
Capitellidae (LPIL)												
<i>Carazziella hobsonae</i>												
<i>Cauleriella sp. J</i>												
<i>Chaetozone</i> (LPIL)												
<i>Chiridotea tuftsi</i>		35	24	29	40	238	10	27	36	48	10	10
Cirratulidae (LPIL)				10	10							
<i>Corophium</i> (LPIL)												
<i>Corophium acutum</i>												
<i>Corophium insidiosum</i>												
<i>Corophium tuberculatum</i>												
<i>Crangon septemspinosa</i>			10		15	10	15					
<i>Crepidula</i> (LPIL)							30					
<i>Crepidula fornicata</i>												
<i>Crepidula plana</i>							26		40			
Cumacea (LPIL)												
Decapoda (LPIL)												
Decapoda Reptantia (LPIL)												
Diastylidae (LPIL)												
<i>Diastylis sculpta</i>												
<i>Diopatra cuprea</i>								10				
<i>Dipolydora commensalis</i>												
<i>Dipolydora socialis</i>									20			
<i>Dispia uncinata</i>		24	68	28	351	15		18	133	30	365	15

**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	Middle	Middle									
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Donax variabilis</i>		30	30			54	518	1538	102	177	14320	1002
<i>Echinarachnius parma</i>												
Echinoidea (LPIL)				10			10				10	
<i>Edotea triloba</i>		10	10	13	10	10	10	15		25		10
<i>Elasmopus levis</i>												
<i>Emerita talpoida</i>			20		10							
<i>Ensis directus</i>			10							10		
<i>Erichthonius rubricornis</i>												
<i>Eusarsiella zostericola</i>								10				
<i>Euspira heros</i>									10			
Gammaridae (LPIL)										10		13
<i>Gammarus</i> (LPIL)		10	15	13				73		40		15
<i>Gammarus annulatus</i>		43	10	15				473		10	190	20
Gastropoda (LPIL)					10							
<i>Glycera</i> (LPIL)							28				10	
<i>Glycera americana</i>												
<i>Glycera capitata</i>												
<i>Glycera dibranchiata</i>										10		
<i>Glycera sp. E</i>												
Glyceridae (LPIL)							40					
<i>Glycinde solitaria</i>												
<i>Haminoea solitaria</i>												
<i>Harmothoe imbricata</i>												
Haustoriidae (LPIL)		15	10	10	26	60	10	18		10	10	
<i>Hemipodus roseus</i>												
<i>Hippomedon sp. C</i>										10		
Hydrobiidae (LPIL)							20					
Hydrozoa (LPIL)												
<i>Hypereteone heteropoda</i>					10							
<i>Idotea balthica</i>											10	
<i>Idotea sp. A</i>												
Idoteidae (LPIL)								10				

**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	Middle	Middle	Middle								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Ilyanassa trivittata</i>												
Isaeidae (LPIL)												
Ischyroceridae (LPIL)			10									
Isopoda (LPIL)		10										
<i>Jassa falcata</i>		10		10		10		10		10	15	
<i>Jassa marmorata</i>						13						
<i>Leucon americanus</i>				10								
<i>Libinia dubia</i>												
Lineidae (LPIL)				10		20	10		10		10	
Lumbrineridae (LPIL)												
Lysianassidae (LPIL)		10						10				10
<i>Macoma</i> (LPIL)												
<i>Macoma yoldiformis</i>				18								
<i>Mactra fragilis</i>											35	
<i>Magelona</i> (LPIL)											10	
<i>Magelona papillicornis</i>		933	1331	167	2595	123	150	15	10	15	30	20
Magelonidae (LPIL)						10						
Majidae (LPIL)												
<i>Mancocuma stellifera</i>						10						
<i>Marenzelleria viridis</i>												
<i>Mediomastus</i> (LPIL)					10						10	
<i>Mediomastus ambiseta</i>												
<i>Mediomastus californiensis</i>												
<i>Melita</i> (LPIL)												
<i>Mulinia lateralis</i>						35					25	
<i>Mya arenaria</i>												
<i>Mysella planulata</i>											40	
Mysidacea (LPIL)			10									
Mysidae (LPIL)			10		10	23	40					
Mytilidae (LPIL)					15							
<i>Mytilus edulis</i>		16		71	45	49	20	143		162	97	27
Naticidae (LPIL)										10		



**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	Middle	Middle	Middle								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Paranaitis speciosa</i>												
Paraonidae (LPIL)				10								
<i>Paraonis fulgens</i>												
<i>Paraprionospio pinnata</i>												
<i>Pectinaria gouldii</i>				10		10						
Pelecypoda (LPIL)		10	20	33								
<i>Petricola pholadiformis</i>												
<i>Pherusa affinis</i>												
Pholadidae (LPIL)												
<i>Photis macrocoxa</i>												
Phoxocephalidae (LPIL)												
<i>Phyllodoce</i> (LPIL)						20						
<i>Phyllodoce arenae</i>		10		10		16						
Phyllodocidae (LPIL)						10						10
<i>Pitar morrhuanus</i>												
Podocopida (LPIL)												
<i>Politilana polita</i>												
<i>Polydora cornuta</i>				10								
<i>Polygordius</i> (LPIL)							10	15		10		
Polynoidae (LPIL)						10						
Portunidae (LPIL)												
<i>Protohaustorius</i> (LPIL)												
<i>Protohaustorius sp. B</i>	18	41	27	39		211	98	79				
<i>Protohaustorius wigleyi</i>									31	257	30	
<i>Psammonyx nobilis</i>	4760	530	20	180	1243	102	115	70	415	275		
<i>Pseudoleptocuma minor</i>			10	10	10		16					10
<i>Pseudunciola obliquua</i>		10									10	
Pyramidellidae (LPIL)												
<i>Rhepoxynius hudsoni</i>					10						10	
Rhynchozoela (LPIL)	40	24	22	34	20	80	46	16	28	35		
<i>Sabellaria vulgaris</i>						81						
<i>Scolecopsis squamata</i>					59	49	23	60	20			



**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	North	North									
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Acanthohaustorius</i> (LPIL)					30							
<i>Acanthohaustorius intermedius</i>												
<i>Acanthohaustorius millsii</i>		30	40	23	78	58	91	116	288	79	226	1221
Acarina (LPIL)			10									
Actiniaria (LPIL)				20				10				
<i>Albunea paretii</i>												
<i>Americamysis bigelowi</i>			20		53		15					
<i>Americhelidium americanum</i>		10	22	13	20	17					30	18
<i>Ampelisca</i> (LPIL)												10
<i>Ampharete</i> (LPIL)			125	30								
<i>Ampharete americana</i>		30	346	20		15						
<i>Ampharete finmarchica</i>				10								
Ampharetidae (LPIL)		20	118	20	34	27	3750	30	10	59		256
Amphipoda (LPIL)							20					
<i>Ampithoe</i> (LPIL)												
<i>Ampithoe longimana</i>												
<i>Ampithoe valida</i>					10							
<i>Anadara ovalis</i>							10					
<i>Ancinus depressus</i>												
Aoridae (LPIL)					30							
<i>Apoprionospio pygmaea</i>												
Archannelida Family A												
Arcidae (LPIL)					10		50					
<i>Aricidea</i> (LPIL)				20					10			
<i>Aricidea catherinae</i>							10					
<i>Asabellides oculata</i>						10	3429	10		123	65	524
Ascidiacea (LPIL)							10					
<i>Astarte castanea</i>							547	15				
Asteroidea (LPIL)							33					
<i>Balanoglossus</i> (LPIL)												
<i>Bathyporeia</i> (LPIL)												
<i>Bathyporeia parkeri</i>		10	10			10					10	

**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	North	North									
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Bathyporeia quoddyensis</i>									10			
Bivalvia (LPIL)						20	50	15		20		
Bodotriidae (LPIL)		10										
<i>Callinectes</i> (LPIL)												
<i>Callinectes similis</i>												
<i>Calyptreaea centralis</i>							50					
Calyptraeidae (LPIL)							51					
<i>Cancer irroratus</i>							15					
<i>Capitella capitata</i>					10					10		
Capitellidae (LPIL)					10							
<i>Carazziella hobsonae</i>												
<i>Caulleriella sp. J</i>												
<i>Chaetozone</i> (LPIL)												
<i>Chiridotea tuftsi</i>		23	180	18	47	162	10	33	70	60	47	110
Cirratulidae (LPIL)				13	10			10	10	10		10
<i>Corophium</i> (LPIL)			10									10
<i>Corophium acutum</i>												10
<i>Corophium insidiosum</i>				20								
<i>Corophium tuberculatum</i>												
<i>Crangon septemspinosa</i>			10		10	10	18		10			
<i>Crepidula</i> (LPIL)												
<i>Crepidula fornicata</i>							50					
<i>Crepidula plana</i>							102					
Cumacea (LPIL)												10
Decapoda (LPIL)												10
Decapoda Reptantia (LPIL)			10									
Diastylidae (LPIL)												
<i>Diastylis sculpta</i>												
<i>Diopatra cuprea</i>							10					
<i>Dipolydora commensalis</i>												
<i>Dipolydora socialis</i>							10					
<i>Dispia uncinata</i>		22	68	62	192	10		20	251	55	840	60





**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	North	North	North								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Neomysis americana</i>			10				50		30			
<i>Neopanope sayi</i>												
Nephtyidae (LPIL)		15	20		18	10	139		15	13		
<i>Nephtys (LPIL)</i>												
<i>Nephtys buccera</i>		10	10	15	10	17	24	10	15	18		43
<i>Nephtys picta</i>			10					23	15	17	15	25
Nereidae (LPIL)												
Nereididae (LPIL)							10		10			
<i>Nereis succinea</i>												
<i>Neverita duplicata</i>												
<i>Nucula proxima</i>							20	10		10		10
<i>Odostomia gibbosa</i>							20					
Oedocerotidae (LPIL)				10								
Oligochaeta (LPIL)				127				23	10	10		
Onuphidae (LPIL)							18					
<i>Onuphis eremita</i>			10								20	
<i>Orbinia americana</i>										10	20	10
Ostraoda (LPIL)												
<i>Ovalipes (LPIL)</i>									10			
<i>Ovalipes ocellatus</i>			20		10		10		40		10	
<i>Ovalipes stephensoni</i>												
<i>Owenia fusiformis</i>												
<i>Oxyurostylis smithi</i>										10	10	17
Paguridae (LPIL)							240					
<i>Pagurus (LPIL)</i>			10			10	33		20		10	10
<i>Pagurus longicarpus</i>						20	10		10			
<i>Pagurus politus</i>			10		10			10		10		
Palaemonidae (LPIL)												
<i>Parahaustorius (LPIL)</i>					10	10						
<i>Parahaustorius attenuatus</i>		10		15	20	17		10		20		10
<i>Parahaustorius longimerus</i>					20	15				10		
<i>Parametopella cypris</i>							10					

**Appendix Table 2-3. Abundance of Nearshore Taxa (Numbers of Animals/m<sup>2</sup>) (Continued)**

Taxa	Area	North	North	North								
	Date	May-95	Sept.- 95	May-96	Sept.- 96	May-97	Sept.- 97	May-98	Sept.- 98	May-99	Sept.- 99	May-00
<i>Paranaitis speciosa</i>												
Paraonidae (LPIL)						10						
<i>Paraonis fulgens</i>			10						10		10	
<i>Paraprionospio pinnata</i>												
<i>Pectinaria gouldii</i>							10				10	
Pelecypoda (LPIL)				43	10							
<i>Petricola pholadiformis</i>										20	10	10
<i>Pherusa affinis</i>							13					
Pholadidae (LPIL)												
<i>Photis macrocoxa</i>				10								
Phoxocephalidae (LPIL)			10									
<i>Phyllodoce</i> (LPIL)												
<i>Phyllodoce arenae</i>			13		15		28					40
Phyllodocidae (LPIL)							60					
<i>Pitar morrhuanus</i>												
Podocopida (LPIL)												10
<i>Politilana polita</i>					15					10		10
<i>Polydora cornuta</i>			10	10								
<i>Polygordius</i> (LPIL)			30					190	10	11	37	
Polynoidae (LPIL)				10			17					
Portunidae (LPIL)			10									
<i>Protohaustorius</i> (LPIL)		20			10							
<i>Protohaustorius sp. B</i>			25	13	37	20	16	10	33			
<i>Protohaustorius wigleyi</i>										47	52	50
<i>Psammonyx nobilis</i>		10			65	127	114	10	10	460	17	
<i>Pseudoleptocuma minor</i>				10	17	12		12	10	10		10
<i>Pseudunciola obliquua</i>			15			10						
Pyramidellidae (LPIL)												
<i>Rhepoxynius hudsoni</i>							10					
Rhynchochoela (LPIL)			20	35	47	18	129	31	26	37	20	23
<i>Sabellaria vulgaris</i>				10		10	270					
<i>Scolecopsis squamata</i>						45	77	26	302	17		10



**Appendix Table 2-4. Analysis of Variance (ANOVA) Results- Biannual Data**

**Abundance**

Source	SS	DF Num	F Ratio	Prob>F
Area	0.2517	2	0.1954	0.8234
Depth	9.0020	2	6.9895	0.0029
Date	5.9236	17	0.5411	0.9105
Area*Depth	9.9736	4	3.8719	0.0068
Area*Date	11.4866	34	0.5246	0.9793
Depth*Date	20.0506	34	0.9158	0.6026
Area*Depth*Date	67.7672	68	1.5476	0.0030
Error	1615.0668	2508		

**Biomass**

Source	SS	DF Num	F Ratio	Prob>F
Area	2.7671	2	0.7845	0.4644
Depth	11.6698	2	3.3086	0.0486
Date	58.9712	17	1.9670	0.0457
Area*Depth	4.4041	4	0.6243	0.6467
Area*Date	0.0063	34	0.0001	1.0000
Depth*Date	10.8594	34	0.1811	1.0000
Area*Depth*Date	539.9615	68	4.5026	<.0001
Error	4422.9992	2508		

**Intertidal Taxa**

Source	SS	DF Num	F Ratio	Prob>F
Area	0.0238	2	0.1680	0.8464
Depth	0.1203	1	0.5860	0.4587
Date	3.9672	12	1.4767	0.2471
Area*Depth	0.3036	2	2.9083	0.0740
Area*Date	1.6974	24	1.3551	0.2311
Depth*Date	2.4642	12	3.9345	0.0021
Area*Depth*Date	1.2526	24	2.1353	0.0011
Error	55.2893	2262		

**Nearshore Taxa**

Source	SS	DF Num	F Ratio	Prob>F
Area	0.6690	2	2.7218	0.0901
Date	4.9318	10	4.0130	0.0040
Area*Date	2.4579	20	4.8294	<.0001
Error	7.5578	297		



**Appendix Table 2-6. Abundance of Monthly Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) from 1999-2000**

Taxa	Area Date	South May-99	South Jun-99	South Jul-99	South Aug-99	South Sep-99	South Oct-99	South Nov-99	South Dec-99	South Jan-00	South Feb-00	South Mar-00	South Apr-00	South May-00
Amphipoda (LPIL)					226									
<i>Amphiporeia virginiana</i>			226											
Aoridae (LPIL)							226							
Ascidiacea (LPIL)														
Bivalvia (LPIL)														
Capitellidae (LPIL)										226				
Cirratulidae (LPIL)														
<i>Crepidula plana</i>									226					
<i>Donax variabilis</i>			678	678	452		226							452
Echinoidea (LPIL)					226									
<i>Emerita talpoida</i>		678		735	226	565	8023	1017	678	678			791	1243
<i>Gammarus</i> (LPIL)			226											
Gastropoda (LPIL)				226										
<i>Haustorius canadensis</i>							452	226	226					
Hippidae (LPIL)														
<i>Jassa falcata</i>														226
<i>Levinsenia gracilis</i>													226	
Lumbrineridae (LPIL)														
<i>Microphthalmus</i> (LPIL)			904											
<i>Microphthalmus aberrans</i>				226										
<i>Microphthalmus sp. G</i>					527	791								
<i>Mytilus edulis</i>		9944	961	603	678			226	226			339	904	527
<i>Nephtys bucera</i>														
Oligochaeta (LPIL)				452			226							
Ostracoda (LPIL)		226												
<i>Ovalipes ocellatus</i>					226									
<i>Parahaustorius attenuatus</i>			226											
<i>Parahaustorius longimerus</i>														
Paraonidae (LPIL)									226					
<i>Petricola pholadiformis</i>								226						
Pleustidae (LPIL)								226						

**Appendix Table 2-6. Abundance of Monthly Intertidal Taxa (Numbers of Animals/m<sup>2</sup>) from 1999-2000 (Continued)**

Taxa	Area	South												
	Date	May-99	Jun-99	Jul-99	Aug-99	Sep-99	Oct-99	Nov-99	Dec-99	Jan-00	Feb-00	Mar-00	Apr-00	May-00
<i>Polygordius</i> (LPIL)							603	226				226		
<i>Protodriloides</i> (LPIL)			2204	2147	490	3119	377	2034	1130	339	1130	678	565	2411
<i>Pseudunciola obliquua</i>							226							
Rhynchocoela (LPIL)		29945	24973	28325	44823	9906	452	3880	3560	1220	1639	7865	6177	37403
<i>Scolelepis squamata</i>		101248	9537	44115	62640	20076	10260	84637	21809	452	12340	2938	1695	34013
Sipuncula (LPIL)							904	1017						
Spionidae (LPIL)		1220	678											
<i>Spisula solidissima</i>								226						
<i>Streblospio benedicti</i>														
<i>Tellina</i> (LPIL)				226										
Tubificidae (LPIL)										226				
<i>Turbonilla interrupta</i>												226		226





**Appendix Table 2-7. Biomass Composition of Monthly Infaunal Samples 1997**

<b>97May</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
Annelid	5.38	94.44	99.72	99.13	99.99	99.59	96.92	35.34	98.47	99.94	99.99	98.07
Crustacea	94.23	-----	-----	-----	-----	-----	-----	64.47	-----	0.03	-----	1.93
Mollusc	0.10	2.78	0.01	-----	-----	0.12	2.84	0.19	0.72	0.03	-----	-----
Echinoderm	-----	-----	-----	0.29	-----	-----	-----	-----	-----	-----	-----	-----
Misc	0.29	2.78	0.27	0.58	-----	0.29	0.24	-----	0.81	-----	0.01	-----
<b>97June</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
Annelid	99.60	99.83	98.72	99.80	98.81	96.28	94.72	99.61	96.38	99.18	97.56	99.89
Crustacea	-----	-----	-----	-----	-----	-----	-----	-----	2.17	-----	0.01	-----
Mollusc	0.39	0.12	-----	0.01	-----	2.11	0.96	0.24	1.07	0.03	1.90	-----
Echinoderm	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Misc	-----	0.05	1.28	0.19	1.19	1.60	4.33	0.15	0.38	0.79	0.53	0.11
<b>97July</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
Annelid	20.00	100.00	33.33	99.46	98.87	98.77	99.26	-----	0.83	99.81	5.59	96.49
Crustacea	40.00	-----	-----	0.41	-----	1.22	-----	99.81	98.35	0.10	94.10	2.89
Mollusc	20.00	-----	33.33	-----	-----	-----	-----	0.19	-----	-----	-----	-----
Echinoderm	0.00	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Misc	20.00	-----	33.33	0.12	1.13	0.01	0.74	-----	0.83	0.10	0.31	0.62
<b>97Aug</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
Annelid	0.01	99.80	1.36	97.45	50.13	99.08	0.99	0.07	78.06	0.08	-----	99.80
Crustacea	94.73	-----	98.49	-----	-----	-----	27.21	84.78	21.86	16.35	99.87	-----
Mollusc	5.25	-----	-----	-----	49.69	-----	71.80	15.07	-----	83.42	-----	-----
Echinoderm	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Misc	0.01	0.20	0.15	2.55	0.18	0.92	0.01	0.07	0.08	0.15	0.13	0.20
<b>97Sept</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
Annelid	12.28	99.28	7.25	99.98	99.31	53.09	99.87	-----	98.74	99.71	99.78	2.16
Crustacea	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	96.38
Mollusc	87.68	-----	92.68	-----	-----	-----	-----	100.00	0.18	-----	-----	0.02
Echinoderm	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Misc	0.04	0.72	0.08	0.02	0.69	46.91	0.13	0.00	1.08	0.29	0.22	1.45

**Appendix Table 2-7. Biomass Composition of Monthly Infaunal Samples 1997**

<b>97Oct</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelid</b>	98.41	99.55	99.44	99.93	99.90	96.43	99.39	98.72	95.28	99.97	99.66	99.94
<b>Crustacea</b>	-----	0.15	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusc</b>	-----	-----	-----	-----	-----	-----	0.01	-----	-----	-----	-----	-----
<b>Echinoderm</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Misc</b>	1.59	0.30	0.56	0.07	0.10	3.57	0.60	1.28	4.72	0.03	0.34	0.06
<b>97Nov</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelid</b>	98.78	10.03	66.37	47.75	-----	31.10	-----	0.22	96.77	45.33	55.74	-----
<b>Crustacea</b>	-----	89.81	33.48	51.25	-----	68.68	-----	99.55	-----	53.33	-----	-----
<b>Mollusc</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	43.93	-----
<b>Echinoderm</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Misc</b>	1.22	0.15	0.15	1.00	-----	0.22	-----	0.22	3.23	1.33	0.33	100.00

**Appendix Table 2-8. Biomass Composition of Monthly Infaunal Samples 1999-2000**

<b>99May</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	15.92	91.64	99.33	97.01	97.39	96.45	70.49	98.36	99.27	92.66	58.73	95.08
<b>Crustacea</b>	82.71	6.71	-----	-----	0.01	-----	28.14	0.70	-----	5.50	40.23	4.60
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	1.35	1.58	0.21	0.39	1.99	3.23	1.15	0.50	0.72	0.59	-----	0.02
<b>Other Taxa</b>	0.02	0.07	0.45	2.60	0.61	0.32	0.23	0.44	0.02	1.25	1.04	0.29
<b>99June</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	86.69	96.66	-----	96.65	49.66	91.24	67.38	97.24	99.82	25.27	40.18	6.33
<b>Crustacea</b>	-----	-----	35.05	-----	0.23	0.09	32.47	0.49	-----	68.95	59.53	93.63
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	8.27	1.82	64.92	1.02	49.66	8.54	0.15	1.45	-----	5.76	0.06	-----
<b>Other Taxa</b>	5.05	1.52	0.03	2.33	0.46	0.13	-----	0.82	0.18	0.02	0.23	0.04
<b>99July</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	88.23	52.89	11.21	69.06	0.04	96.23	99.96	99.23	99.93	99.86	99.88	31.36
<b>Crustacea</b>	-----	47.05	87.51	30.34	99.89	-----	-----	-----	-----	-----	-----	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	11.68	-----	0.35	-----	-----	2.52	-----	0.70	-----	-----	-----	68.34
<b>Other Taxa</b>	0.09	0.06	0.94	0.60	0.06	1.25	0.04	0.07	0.07	0.14	0.12	0.30
<b>99Aug</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	69.58	47.45	76.97	98.78	88.02	99.89	99.50	97.63	2.96	99.67	0.56	-----
<b>Crustacea</b>	-----	-----	0.51	-----	11.77	-----	-----	-----	71.85	-----	99.06	35.39
<b>Echinodermata</b>	1.13	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	27.21	49.63	21.09	-----	-----	-----	-----	-----	23.62	-----	-----	64.60
<b>Other Taxa</b>	2.08	2.92	1.43	1.22	0.22	0.11	0.50	2.37	1.57	0.33	0.37	-----
<b>99Sept</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	60.31	88.53	93.81	99.25	21.78	99.68	83.40	99.74	99.24	59.67	15.18	25.21
<b>Crustacea</b>	36.94	-----	-----	-----	77.45	-----	15.59	-----	-----	39.44	79.27	50.99
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	-----	-----	-----	-----	-----	-----	0.19	-----	-----	-----	-----	-----
<b>Other Taxa</b>	2.75	11.47	6.19	0.75	0.76	0.32	0.83	0.26	0.76	0.89	5.55	23.80

**Appendix Table 2-8. Biomass Composition of Monthly Infaunal Samples 1999-2000**

<b>99Oct</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	87.92	32.74	1.51	8.18	0.75	9.84	97.46	6.84	99.69	97.87	17.41	100.00
<b>Crustacea</b>	12.05	66.45	98.23	91.82	99.25	90.16	-----	73.74	-----	-----	70.49	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	-----	-----	0.04	-----	-----	-----	2.48	19.41	-----	1.65	-----	-----
<b>Other Taxa</b>	0.03	0.81	0.22	-----	-----	-----	0.06	0.00	0.31	0.48	12.10	-----
<b>99Nov</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	18.95	99.22	99.48	99.96	99.94	99.38	80.63	100.00	100.00	91.63	-----	73.54
<b>Crustacea</b>	75.38	0.59	0.26	-----	-----	0.35	-----	-----	-----	-----	96.74	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	3.56	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.87	-----
<b>Other Taxa</b>	2.11	0.19	0.26	0.03	0.06	0.27	19.37	-----	-----	8.37	2.39	26.46
<b>99Dec</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	77.54	98.69	93.50	7.99	98.39	-----	22.16	-----	-----	-----	100.00	-----
<b>Crustacea</b>	-----	-----	6.16	90.38	1.38	100.00	-----	-----	-----	-----	-----	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	17.85	-----	-----	1.63	-----	-----	-----	100.00	-----	-----	-----	-----
<b>Other Taxa</b>	4.62	1.31	0.33	-----	0.22	-----	77.84	-----	-----	-----	-----	100.00
<b>00Jan</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	49.43	100.00	86.91	3.57	10.65	32.43	100.00	-----	-----	-----	-----	-----
<b>Crustacea</b>	-----	-----	-----	91.64	88.26	-----	-----	-----	-----	-----	-----	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Other Taxa</b>	50.57	-----	13.09	4.79	1.09	67.57	-----	100.00	-----	-----	-----	-----
<b>00Feb</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	-----	99.44	97.22	4.83	100.00	100.00	99.53	100.00	99.73	99.09	-----	95.96
<b>Crustacea</b>	-----	-----	-----	95.17	-----	-----	-----	-----	-----	-----	-----	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	3.59
<b>Other Taxa</b>	100.00	0.56	2.78	-----	-----	-----	0.47	-----	0.27	0.91	100.00	0.45

**Appendix Table 2-8. Biomass Composition of Monthly Infaunal Samples 1999-2000**

<b>00Mar</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	99.72	98.35	99.84	97.89	97.36	82.29	99.96	48.24	99.48	98.08	98.53	96.86
<b>Crustacea</b>	-----	-----	-----	-----	-----	-----	-----	51.60	-----	-----	-----	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	0.07	0.54	-----	-----	2.55	13.67	-----	-----	-----	1.26	1.47	-----
<b>Other Taxa</b>	0.21	1.11	0.16	2.11	0.10	4.04	0.04	0.16	0.52	0.66	-----	3.14
<b>00Apr</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	31.82	94.19	99.75	60.80	8.71	-----	91.30	99.93	13.45	99.08	99.60	99.85
<b>Crustacea</b>	-----	-----	-----	39.14	91.26	-----	-----	-----	86.04	-----	-----	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	-----	-----	-----	-----	-----	74.13	-----	-----	-----	-----	-----	0.03
<b>Other Taxa</b>	68.18	5.81	0.25	0.06	0.03	25.87	8.70	0.07	0.51	0.92	0.40	0.11
<b>00May</b>	<b>Station 1</b>	<b>Station 2</b>	<b>Station 3</b>	<b>Station 4</b>	<b>Station 5</b>	<b>Station 6</b>	<b>Station 21</b>	<b>Station 22</b>	<b>Station 23</b>	<b>Station 24</b>	<b>Station 25</b>	<b>Station 26</b>
<b>Annelida</b>	99.42	99.88	99.83	96.56	50.10	59.69	99.82	83.17	99.90	95.50	64.04	98.77
<b>Crustacea</b>	-----	-----	-----	0.35	49.12	40.15	-----	-----	-----	-----	35.81	-----
<b>Echinodermata</b>	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
<b>Mollusca</b>	0.02	-----	0.11	0.44	0.57	-----	-----	6.93	-----	4.05	0.06	-----
<b>Other Taxa</b>	0.56	0.11	0.06	2.65	0.21	0.16	0.18	9.90	0.10	0.45	0.09	1.23

**Table 2-2. Relative Abundance and Occurrence of Dominant Intertidal Taxa**

Taxa	Area	Total		South		Middle		North	
		Abund.	Occur.	Abund.	Occur.	Abund.	Occur.	Abund.	Occur.
Rhynchocoela (LPIL)		51.44	35.64	44.27	36.15	66.24	36.03	36.55	34.74
<i>Scolelepis squamata</i>		34.87	30.00	45.35	30.38	26.80	30.26	35.11	29.36
Oligochaeta (LPIL)		8.32	16.45	4.62	12.95	4.37	18.21	19.01	18.21
<b>Mytilus edulis</b>		1.60	14.15	2.38	13.97	0.82	12.82	1.89	15.64
<i>Protodriloides</i> (LPIL)		0.82	6.07	1.26	6.03	0.37	6.03	1.02	6.15
<i>Corophium tuberculatum</i>		0.45	0.17	---	---	<0.01	0.13	1.69	0.38
<i>Microphthalmus</i> (LPIL)		0.42	2.99	0.04	1.15	0.24	2.82	1.17	5.00
<i>Emerita talpoida</i>		0.38	4.40	0.35	6.41	0.11	2.56	0.84	4.23
<i>Microphthalmus</i> sp. G		0.36	2.09	0.15	2.44	0.07	1.54	1.08	2.31
<i>Microphthalmus aberrans</i>		0.22	0.64	0.06	0.90	0.41	1.41	0.20	0.98
<i>Polygordius</i> (LPIL)		0.16	0.60	0.49	1.15	0.01	0.38	0.01	0.26
<b>Spionidae (LPIL)</b>		0.12	2.82	0.03	1.41	0.09	1.67	0.07	1.97
<i>Parahaustorius attentuatus</i>		0.08	2.95	0.05	2.44	0.13	2.18	0.08	2.52
<i>Donax variabilis</i>		0.04	1.15	0.07	1.15	0.02	0.64	0.05	0.98
<i>Ovalipes ocellatus</i>		0.01	0.77	0.03	1.79	0.09	1.67	0.04	1.41
<i>Parahaustorius longimerus</i>		0.01	0.64	0.03	1.28	0.03	0.77	0.02	0.90
<i>Haustorius</i> sp. C		<0.01	0.26	0.03	1.28	0.02	0.51	0.02	0.68
Pisionidae (LPIL)		---	---	<0.01	0.13	0.25	1.03	0.07	0.38

\*Taxa in **bold** excluded from final analyses (see text)

**Table 2-3. Relative Abundance and Occurrence of Dominant Nearshore Infauna**

Taxa	Area	Total		South		Middle		North	
		Abund.	Occur.	Abund	Occur	Abund	Occur	Abund	Occur
<i>Donax variabilis</i>		74.84	46.97	77.90	47.27	48.56	50.00	78.85	43.64
<i>Magelona papillicornis</i>		5.75	60.00	5.40	63.64	13.79	60.91	3.99	55.45
<i>Tellina agilis</i>		3.60	100.00	4.12	100.00	5.13	98.18	2.73	100.00
<i>Spisula solidissima</i>		2.11	64.85	1.11	61.82	5.19	67.27	2.23	65.45
<i>Acanthohaustorius millsii</i>		2.03	80.61	2.15	89.09	3.81	80.91	1.45	71.82
<i>Psammonyx nobilis</i>		1.94	36.36	2.37	57.27	6.77	30.00	0.29	21.82
<b>Ampharetidae (LPIL)</b>		1.88	37.58	0.27	30.91	3.40	37.27	2.97	44.55
<i>Asabellides oculata</i>		1.33	26.06	0.65	26.36	0.69	21.82	2.13	30.00
<i>Dispio uncinata</i>		0.98	59.09	0.46	55.45	2.62	66.36	1.03	55.45
<b><i>Mytilus edulis</i></b>		0.92	53.64	0.56	56.36	1.26	46.36	1.15	58.18
<i>Chiridotea tuftsi</i>		0.55	61.21	0.59	66.36	1.15	60.00	0.36	57.27
<i>Protohaustorius sp. B</i>		0.47	41.52	0.76	60.00	1.16	43.64	0.04	20.91
<i>Gammarus annulatus</i>		0.40	21.21	0.45	21.82	0.84	19.09	0.25	22.73
<i>Protohaustorius wigleyi</i>		0.36	16.36	0.59	20.91	0.81	18.18	0.04	10.00
<i>Astarte castanea</i>		0.29	8.18	0.16	5.45	0.33	8.18	0.40	10.91
<i>Scolecopsis squamata</i>		0.23	20.30	0.17	13.64	0.31	20.91	0.27	26.36
Rhynchocoela (LPIL)		0.22	58.79	0.11	60.91	0.68	57.27	0.20	58.18
<i>Spiophanes bombyx</i>		0.11	26.67	0.16	33.64	0.22	24.55	0.04	21.82
<i>Ampharete americana</i>		0.08	10.00	0.04	15.45	0.06	5.45	0.13	9.09
<i>Nephtys buccera</i>		0.06	27.27	0.07	30.91	0.13	23.64	0.04	27.27
<i>Neomysis americana</i>		0.06	8.48	0.10	8.18	0.06	7.27	0.03	10.00
<i>Parahaustorius attenuatus</i>		0.05	14.55	0.07	17.27	0.14	11.82	0.02	14.55
<i>Nephtys picta</i>		0.05	19.70	0.05	21.82	0.08	14.55	0.03	22.73
<i>Edotea triloba</i>		0.04	16.06	0.06	21.82	0.06	13.64	0.02	12.73
<i>Americhelidium americanum</i>		0.03	19.70	0.03	24.55	0.05	13.64	0.03	20.91
<i>Pagurus</i> (LPIL)		0.03	11.52	0.03	12.73	0.04	7.27	0.03	14.55
<i>Americamysis bigelowi</i>		0.03	8.79	0.02	6.36	0.08	10.91	0.02	9.09
<i>Phyllodoce arenae</i>		0.03	11.82	0.04	17.27	0.03	7.27	0.02	10.91
<i>Polygordius</i> (LPIL)		0.03	10.00	0.02	10.91	0.02	6.36	0.03	12.73
<i>Pseudoleptocuma minor</i>		0.02	15.45	0.02	11.82	0.05	12.73	0.02	21.82

\*Taxa in **bold** excluded from final analyses (see text)

**Table 2-4. Analysis of Variance (ANOVA) Results- Biannual Data\***

<b>MLW Abundance</b>				
Source	SS	DF	F Ratio	Prob>F
Area	3.1826	2	2.5465	0.0993
Date	3.4631	12	0.4618	0.9178
Area*Date	94.5518	24	6.3045	<.0001
Error	706.7598	1131		

<b>MLW -1m Abundance</b>				
Source	SS	DF	F Ratio	Prob>F
Area	1.1604	2	0.7558	0.4805
Date	7.7220	12	0.8383	0.6134
Area*Date	48.1442	24	2.6131	<.0001
Error	868.2349	1131		

<b>Nearshore Abundance</b>				
Source	SS	DF	F Ratio	Prob>F
Area	1.3305	2	3.2160	0.0615
Date	11.7774	10	5.6939	0.0005
Area*Date	13.4224	20	3.2446	<.0001
Error	61.4319	297		

<b>MLW Biomass</b>				
Source	SS	DF	F Ratio	Prob>F
Area	4.6118	2	1.2794	0.2965
Date	55.3450	12	2.5589	0.0242
Area*Date	256.4352	24	5.9281	<.0001
Error	2038.5013	1131		

<b>MLW-1m Biomass</b>				
Source	SS	DF	F Ratio	Prob>F
Area	0.4076	2	0.0977	0.9073
Date	50.5510	12	2.0194	0.0690
Area*Date	1998.2297	24	3.9594	<.0001
Error	2359.3130	1131		

<b>Nearshore Biomass</b>				
Source	SS	DF	F Ratio	Prob>F
Area	2.4063	2	2.5778	0.1009
Date	11.0007	10	2.3584	0.0491
Area*Date	28.2761	20	3.0292	<.0001
Error	138.6197	297		

<b>MLW Taxa</b>				
Source	SS	DF	F Ratio	Prob>F
Area	0.5899	2	0.2429	0.7862
Date	9.0502	12	0.6211	0.8038
Area*Date	77.3406	24	2.6539	<.0001
Error	1373.3420	1131		

<b>MLW-1m Taxa</b>				
Source	SS	DF	F Ratio	Prob>F
Area	5.0113	2	1.9086	0.1701
Date	70.4679	12	4.4732	0.0009
Area*Date	65.5492	24	2.0805	<.0001
Error	1484.7667	1131		

<b>Nearshore Taxa</b>				
Source	SS	DF	F Ratio	Prob>F
Area	148.5077	2	3.5970	0.0463
Date	1434.5596	10	6.9498	0.0001
Area*Date	912.7890	20	2.2110	<.0024
Error	6130.6000	297		

**Table 2-5. Correlations Between Taxa and MDS Axes\***

Taxon	Axis	
	1	2
<i>Ampelisca abdita</i>	-0.010	-0.265
Archiannelida (LPIL)	-0.027	-0.240
<i>Corophium tuberculatum</i>	0.021	-0.120
<i>Donax variabilis</i>	<b>-0.402</b>	<b>0.790</b>
<i>Emerita talpoida</i>	0.373	-0.284
Hirudinea (LPIL)	0.100	-0.238
<i>Magelona papilicornis</i>	<b>-0.920</b>	<b>0.622</b>
<i>Mediomastus ambiseta</i>	-0.039	-0.237
<i>Microphthalmus aberrans</i>	0.122	<b>-0.439</b>
<i>Microphthalmus hartmanae</i>	0.055	-0.164
<i>Microphthalmus sp. G</i>	0.350	-0.285
<i>Microphthalmus listensis</i>	0.162	-0.130
Oligochaeta (LPIL)	<b>0.516</b>	<b>-0.811</b>
<i>Parahaustorius longimerus</i>	0.023	-0.078
<i>Parahaustorius attentuatus</i>	0.063	-0.114
<i>Polygordius</i> (LPIL)	-0.102	0.103
<i>Protodriloides</i> (LPIL)	0.367	-0.094
Rhynchocoela (LPIL)	<b>0.903</b>	<b>-0.836</b>
<i>Scolelepis squamata</i>	<b>0.930</b>	<b>-0.740</b>

\*Values in bold considered to be significant correlations

**Table 2-6. Analysis of Variance (ANOVA) Results - Monthly Data - 1997**

**Abundance**

Source	SS	DF Num	F Ratio	Prob>F
Area	0.0974	1	0.1047	0.7572
Date	3.8805	6	0.6954	0.6649
Area*Date	14.8796	6	2.6664	0.0160
Error	221.3548	238		

**Biomass**

Source	SS	DF Num	F Ratio	Prob>F	Power
Area	0.8438	1	0.3453	0.5782	0.0507
Date	60.0846	6	4.0975	0.0550	0.0616
Area*Date	18.4138	6	1.2557	0.2788	0.4897
Error	581.6682	238			

**Taxa Richness**

Source	SS	DF Num	F Ratio	Prob>F	Power
Area	0.0003	1	0.0128	0.9137	0.0500
Date	0.1583	6	1.0855	0.4616	0.0530
Area*Date	0.2138	6	1.4662	0.1905	0.5646
Error	5.7861	238			

**Table 2-7. Analysis of Variance (ANOVA) Results - Monthly Data 1999-2000**

**Abundance**

<b>Source</b>	<b>SS</b>	<b>DF Num</b>	<b>F Ratio</b>	<b>Prob&gt;F</b>
Area	5.5938	1	7.3749	0.0188
Date	30.3031	12	3.3293	0.0236
Area*Date	67.9997	12	7.4709	<0.0001
Error	335.2552	442		

**Biomass**

<b>Source</b>	<b>SS</b>	<b>DF Num</b>	<b>F Ratio</b>	<b>Prob&gt;F</b>
Area	8.9809	1	4.3961	0.5790
Date	71.4768	12	2.9254	0.0375
Area*Date	179.2814	12	7.3377	<0.0001
Error	899.9482	442		

**Taxa Richness**

<b>Source</b>	<b>SS</b>	<b>DF Num</b>	<b>F Ratio</b>	<b>Prob&gt;F</b>
Area	0.2184	1	9.6768	0.0090
Date	1.7601	12	6.5001	0.0014
Area*Date	0.9594	12	3.5432	<0.0001
Error	9.9738	442		

**Table 2-8. Benthic Recovery Rate Estimates**

<b>Area</b>	<b>Year</b>	<b>Parameter</b>	<b>Regression Line*</b>	<b>Time to Recovery (Days)</b>	<b>p Value</b>	<b>r<sup>2</sup> value</b>
South	1997	Abundance	$X = 2.85207 + 0.00301Y$	49	0.0059	0.63
South	1997	Biomass	$X = 2.86079 + 0.00352Y$	38	0.0145	0.55
South	1997	Taxa Richness	$X = 2.89811 + 0.00448Y$	23	0.0072	0.63
Middle	1999	Abundance	$X = 1.60700 + 0.00736Y$	189	0.0026	0.24
Middle	1999	Biomass	$X = 1.13418 + 0.01056Y$	176	0.0016	0.26
Middle	1999	Taxa Richness	$X = 2.47089 + 0.00316Y$	178	0.0002	0.36

\* X = ArcSine-Square Root of % Recovery (100% = 3) and Y = days

**Table 2-9. Representative Infaunal Abundances for Atlantic Coast Sandy Beaches.**

<b>Reference</b>	<b>Site</b>	<b>Abundance/m<sup>2</sup></b>
Larsen and Doggett (1990)	Southern Maine	0-25,000
Sameoto (1969)	Cape Cod, MA	3,000
McDermott (1983)	Avalon Beach, NJ	42,000-86,000
This Study	Asbury Park - Manasquan Inlet , NJ	9,531-14,773
Matta (1977)	Duck, NC	30-800
Diaz and DeAlteris (1982)	Duck, NC	2,300
Dexter (1969)	Morehead City, NC	60
Van Dolah et al. (1994)	Folly Beach, SC	1,000-22,000
Jutta et al. (1999ab)	Myrtle Beach, SC	1,000-25,000
Knott et al. (1983)	Murrells Inlet, SC	1,500-3,200
Gorzelany and Nelson (1987)	Melbourne Beach, FL	1,000-4,200

**Table 2-10. Summary of Beach Nourishment Recovery Rate Estimates**

<b>Reference</b>	<b>Site</b>	<b>Recovery rate (months)</b>
Reilly and Bellis (1983)	Bogue Banks, NC	>12 <sup>sc</sup>
Van Dolah et al. (1994)	Folly Beach, SC	2-3
Jutta et al. (1999ab)	Myrtle Beach, SC	6
Gorzelany and Nelson (1987)	Melbourne Beach, FL	<1*
Salomon and Naughton (1984)	Panama City, FL	2
Rakocinski et al. (1996)	Perdido Bay, FL	>12 <sup>sc</sup>

<sup>sc</sup>Substantial amounts of silts and clays present

\*No impact detected




Figure 2-2. Intertidal and Nearshore benthos sampling locations in the South Area.




Figure 2-3. Intertidal and Nearshore benthos sampling locations in the Middle Area.







Figure 2-4. Intertidal and Nearshore benthos sampling locations in the North Area.




**Figure 2-5. Mean Low Water Abundance (No. Animals/m<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.




**Figure 2-6. Mean Low Water -1m Abundance (No. Animals/m<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.




**Figure 2-7. Nearshore Abundance (No. Animals/m<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.




**Figure 2-8. Mean Low Water Biomass (Grams/m<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.




**Figure 2-9. Mean Low Water -1m Biomass (Grams/m<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant (p<0.0125) linear contrast between means.




**Figure 2-10. Nearshore Biomass (Grams/m<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.



**Figure 2-11. Mean Low Water Taxa Richness (Taxa/44cm<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.



**Figure 2-12. Mean Low Water -1m Taxa Richness (Taxa/44cm<sup>2</sup> ± SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.



**Figure 2-13. Nearshore Taxa Richness (Taxa/0.1m<sup>2</sup>±SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Red arrow indicates South Area nourished; Black arrow indicates Middle Area nourished; Bar indicates a significant ( $p < 0.0125$ ) linear contrast between means.




Figure 2-14. MLW Benthos Biomass Composition- South Area. Arrow indicates when nourishment occurred.




Figure 2-15. MLW Benthos Biomass Composition- Middle Area. Arrow indicates when nourishment occurred.






Figure 2-16. MLW Benthos Biomass Composition- North Area.



**Figure 2-17. MLW-1m Benthos Biomass Composition- South Area.** Arrow indicates when nourishment occurred.



**Figure 2-18. MLW-1m Benthos Biomass Composition- Middle Area.** Arrow indicates when nourishment occurred.






Figure 2-19. MLW-1m Benthos Biomass Composition- North Area.



**Figure 2-20. Nearshore Benthos Biomass Composition- South Area.** Arrow indicates when nourishment occurred.



**Figure 2-21. Nearshore Benthos Biomass Composition- Middle Area.** Arrow indicates when nourishment occurred.






Figure 2-22. Nearshore Benthos Biomass Composition- North Area.

### Intertidal-Nearshore MDS




**Figure 2-23. Intertidal-Nearshore Infaunal NMDS Results.** Black = South Area, Red = Middle Area, Green = North Area; Triangle = MLW, Circle = MLW-1m. Square = Nearshore. Stress = 0.13.


### Intertidal-Nearshore MDS




**Figure 2-24. NMDS Results from Intertidal-Nearshore Samples During and Immediately After Nourishment.** Nearshore Data: S= South, M= Middle, N=North; 7=Sept 1997, 8=May 1998, 9 = Sept. 1999, 0=May 2000. Intertidal Data: S = South, 1<sup>st</sup> M = Middle, N= North, 2<sup>nd</sup> M= MLW, L= MLW-1m; 7=Sept 1997, 8=May 1998, 9 = Sept. 1999, 0=May 2000. Stress = 0.13.



**Figure 2-25. Mean Low Water Mean Grain Size (mm<sub>±SE</sub>) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Arrows indicate when nourishment occurred (1<sup>st</sup> = South, 2<sup>nd</sup> = Middle).



**Figure 2-26. Mean Low Water –1m Mean Grain Size (mm $\pm$ SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Arrows indicate when nourishment occurred (1<sup>st</sup> = South, 2<sup>nd</sup> = Middle).



**Figure 2-27. Nearshore Mean Grain Size (mm $\pm$ SE) by Date.** Black Circles = South Area, Red Circles = Middle Area, Green Triangles = North Area; Arrows indicate when nourishment occurred (1<sup>st</sup> = South, 2<sup>nd</sup> = Middle).





Figure 2-28. MLW Sediment Composition- South Area. Arrow indicates when nourishment occurred.



**Figure 2-29. MLW Sediment Composition- Middle Area.** Arrow indicates when nourishment occurred.




Figure 2-30. MLW Sediment Composition- North Area.




Figure 2-31. MLW-1m Sediment Composition- South Area. Arrow indicates when nourishment occurred.




Figure 2-32. MLW-1m Sediment Composition- Middle Area. Arrow indicates when nourishment occurred.




Figure 2-33. MLW-1m Sediment Composition- North Area.





Figure 2-34. Nearshore Sediment Composition- South Area. Arrow indicates when nourishment occurred.



**Figure 2-35. Nearshore Sediment Composition- Middle Area.** Arrow indicates when nourishment occurred.







Figure 2-36. Nearshore Sediment Composition- North Area.




**Figure 2-37. Monthly Abundance (No. Animals/m<sup>2</sup>±SE) for 1997.** Red Circles = South Area, Black Circles = Middle Area, Red line indicates time during which South Area MLW Stations 1-6 were nourished; Bar indicates a significant ( $p < 0.0167$ ) linear contrast between means.




**Figure 2-38. Monthly Abundance (No. Animals/m<sup>2</sup>±SE) for 1999.** Red Circles = South Area, Black Circles = Middle Area, Red line indicates time during which Middle Area MLW Stations 21-26 were nourished; Bar indicates a significant ( $p < 0.0167$ ) linear contrast between means.




**Figure 2-39. Monthly Biomass (Grams/m<sup>2</sup>+SE) for 1997.** Red Circles = South Area, Black Circles = Middle Area, Red line indicates time during which South Area MLW Stations 1-6 were nourished; Bar indicates a significant ( $p < 0.0167$ ) linear contrast between means.




**Figure 2-40. Monthly Biomass (Grams/m<sup>2</sup>±SE) for 1999.** Red Circles = South Area, Black Circles = Middle Area, Red line indicates time during which Middle Area MLW Stations 21-26 were nourished; Bar indicates a significant ( $p < 0.0167$ ) linear contrast between means.




**Figure 2-41. Monthly Taxa Richness (Taxa/44cm<sup>2</sup> ± SE) for 1997.** Red Circles = South Area, Black Circles = Middle Area, Red line indicates time during which South Area MLW Stations 1-6 were nourished; Bar indicates a significant ( $p < 0.0167$ ) linear contrast between means.




**Figure 2-42. Monthly Taxa Richness (Taxa/44cm<sup>2</sup>±SE) for 1999.** Red Circles = South Area, Black Circles = Middle Area, Red line indicates time during which Middle Area MLW Stations 21-26 were nourished; Bar indicates a significant (p<0.0167) linear contrast between means.




**Figure 2-43. Monthly Abundance (Animals/m<sup>2</sup> ± SE) of *Scololepis squamata*: 1999-2000.** Orange = South Area (reference), Brown = Middle Area (nourished).




**Figure 2-44. Monthly Sediment Mean Grain Size ( $\pm$ SE) for South Area 1997.**  
 Numbers = MLW Stations.




**Figure 2-45. Monthly Sediment Mean Grain Size ( $\pm$ SE) for Middle Area 1997.**  
 Numbers = MLW Stations.




**Figure 2-46. Monthly Sediment Mean Grain Size ( $\pm$ SE) for South Area 1999.**  
 Numbers = MLW Stations.




**Figure 2-47. Monthly Sediment Mean Grain Size ( $\pm$ SE) for Middle Area 1999.**  
 Numbers = MLW Stations.




**Figure 2-48. Surfzone and Nearshore Bottom Water Temperature (Mean C° ±SE).**  
 Filled symbols = Surfzone, Open symbols = Nearshore Bottom water; Circle = South Area, Inverted Triangle = Middle Area, Square = North Area.



**Figure 2-49. Surfzone and Nearshore Bottom Water Salinity (Mean ppt  $\pm$ SE).**  
 Filled symbols = Surfzone, Open symbols = Nearshore Bottom water; Circle = South Area, Inverted Triangle = Middle Area, Square = North Area.



**Figure 2-50. Surfzone and Nearshore Bottom Water Dissolved Oxygen Concentrations (Mean mg/l<sub>±</sub>SE).** Filled symbols = Surfzone, Open symbols = Nearshore Bottom water; Circle = South Area, Inverted Triangle = Middle Area, Square = North Area.



**Figure 2-51. Surfzone and Nearshore Bottom Water pH (Mean<sub>±</sub>SE).** Filled symbols = Surfzone, Open symbols = Nearshore Bottom water; Circle = South Area, Inverted Triangle = Middle Area, Square = North Area.

Likewise, Rakocinski et al. (1996) found delayed recovery of infaunal assemblages at a nourished beach and shallow inshore habitats in Perdido Key, Florida where there were substantial amounts of silts and clays in the fill material. Peterson et al. (2000) have examined impacts of beach nourishment to large beach infauna (*Emerita*, *Donax* and the ghost crab *Ocypode*) at Bogue Sound, North Carolina. The presence of a substantial amount of silts and clays as well as shell hash in the fill materials resulted in infaunal abundances still being reduced 2.5 months after nourishment.

## Methods

**Monitoring Plan and Overall Execution:** Sampling of intertidal benthos took place in two phases. In the first phase, samples were collected twice a year (spring and fall) at 10 sites within each of three areas and at two depths: 2 seasons x 3 areas/season x 10 sites/area x 2 stations/site x 3 cores/station for a total of 360 samples/year. Target areas were South Area, Middle Area, and North Area (Figure 2-1); and station depths were Mean Low Water (MLW) and MLW-1m (Figures 2-2 to 2-4). Station locations for MLW are provided in Appendix Table 2-1.

Replicate samples were collected at 1-2 m intervals with 7.5 cm PVC coring tubes to a depth of 10-15 cm, and a 0.5 mm sieve was used to separate infauna from the sediment. The monitoring plan also called for 120 sediment samples (one per infauna station per season) taken with a 5 cm, PVC coring tube to a depth of 10-15 cm, which were processed for grain-size distribution. Sampling began in June 1994 and was completed in May 2000. The second phase was initiated only when nourishment operations began. In 1997 the first six stations of both the nourished beach (South Area) and one of the reference areas (Middle Area) were sampled at MLW in June, July, August, October, and November (1997). In 1999-2000, when the Middle Area was nourished, the same stations were sampled, but the South Area acted as reference. As will be discussed later, the South Area had recovered sufficiently to serve as a reliable reference site. Samples were taken on a monthly schedule between June 1999 and April 2000. When combined with data from the May and September collections for the same stations, these samples permit a detailed temporal analysis of potential impacts and recovery ranging over a period of seven (1997) to thirteen (1999-2000) months.

Although sampling of nearshore benthos was not part of the original study design, this component was added in 1995 to extend characterization efforts far enough off the beach to capture potential impacts related to changes in profiles resulting from filling operations. Nearshore stations were located along the pre-construction 5.0-6.5 m depth contour, which generally lies just seaward of the terminal points of the existing groin field (Figures 2-2 to 2-4). Characterization of the nearshore benthos also enhanced interpretation of surf zone fish food habits data. Nearshore samples were collected with a single Smith-McIntyre grab (0.1 m<sup>2</sup>) sample directly seaward of each intertidal site/station set, yielding a total of 30 samples per sampling date (3 areas x 10 sites/area x 1 sample/site). Each site was sampled twice a year (May and September) beginning in

1995 and ending in May 2000. A sample for sediment texture analysis was also taken at each sampling point.

The monitoring plan was executed as planned for all years (1994-2000) except for the loss of a few samples in May 1994, when sediment texture samples were lost in the field from site 25 (MLW of the fifth transect at Middle area), and all samples from September 1995 collections. Infauna and sediment samples from Middle area Stations 23 and 33 were also misplaced in September 1998. While the sample losses were unfortunate, the effect on the monitoring program was minimal. Correlations between sediment type and infauna abundance are either non-significant or weak for the remaining samples. Thus, interpretations of the infauna data are not impaired by the decrease in sediment data. Otherwise, all infaunal and sediment samples were taken on schedule for both phases (biannual and monthly) of the project.

***Field and Laboratory Methods:*** The biannual samples (Phase 1) were collected during daytime, spring low tides during May and September of each year. The time of sampling was standardized to permit the most efficient use of field time and to reduce the influence of tidal migrations on taxa abundances (Jones et al. 1998).

Two to three days were required to collect all the samples during each sampling event. The groins were used to guide the distribution of the 10 sites in each sample area, with the goal being to sample the entire area as evenly as practicable. The South Area, which is between Philadelphia Blvd. and Remsen Ave., has 13 groins, which yields 12 inter-groin spaces (Figure 2-2). One sampling site was located near the center of each space, except for the two spaces near the storm-water outlet from Wreck Pond, which were excluded to avoid potential influences from the discharge. This sampling area encompasses 3.7 km of coastline, and the typical distance between transects is 300 m. The Middle Area, which is between Washington Ave. and Seaview Ave., has 12 groins, which yields 11 inter-groin spaces (Figure 2-3). One site was located near the center of each of these 11 spaces, except for the one space near the storm-water outlet from Fletcher Lake, which was excluded to avoid potential influences from the discharge. The sampling area covers 3.3 km of coastline, and the typical distance between sites is 300 m. The North sampling area, which covers 3.4 km between Cedar Ave. and Roosevelt Ave., has 12 groins, which yields 11 inter-groin spaces, but sampling in this area was not as straightforward as in the others (Figure 2-4). The portion near Neptune Ave. (2 inter-groin spaces) could not be sampled because there was no sandy beach during low tide. Furthermore, the groins are not distributed as evenly nor area as uniform in size as in the other areas. As a result, it was necessary to put two sites in two of the inter-groin spaces to provide 10 sites in this area. Distances between sites ranged from 100-220 m. Sampling occurred near the center of each inter-groin space to minimize the influence that groins might have on the immediately adjacent benthic community. In some cases, however, sites were moved away from the center to avoid derelict groins or storm-water outfalls.

Stations were initially located within each area based on standard land survey methods. A base station, usually in the South Area, was established using a transit, Philadelphia rod, and maps of street elevations (height above MLW) prepared by Coastal Planning and Engineering,

Inc. The elevation of the boardwalk above MLW was determined using the transit, rod, and elevation of the street as a reference. The elevation of the beach at the base of the boardwalk was then determined by subtracting the elevation of the boardwalk from the height of the boardwalk above the beach. The transit and rod were then used to locate MLW along the beach face, and the location was marked for that day with a stake. The depth of the water at the stake was used to locate MLW at other locations (i.e., if water depth was 10 cm at the stake at a given time, we assumed other 10 cm deep locations in the study area were also MLW). The sampling crew communicated with an observer at the base station via a cellular phone to obtain information about the depth of water relative to MLW. After several years experience with this method, it was noted that equal levels of accuracy could be obtained basing station locations on beach structure and depth relative to time and tidal height. Specifically, it was found that a small beach step was present at all sites at the Mean Low Water (MLW) mark. Beach steps are common on coarse sandy beaches; the crest of the step is generally associated with the Mean Low Water level (Bauer and Allen, 1995). Using this structure to site the MLW station, the MLW-1m station could be easily determined by depth relative to the depth of water over the MLW station. Thus, if there were 10 cm of water over the MLW station, the MLW-1m station was located at depth of 110 cm. When this procedure was tested against the transect level method in Spring and Fall 1996, no substantive differences could be detected between the two methods.

Infaunal samples were preserved in the field with buffered 10% formalin. In the laboratory at Barry A. Vittor & Associates, Inc. (BVA), Mobile, Alabama, samples were stained with 1% Rose Bengal and transferred to 70% isopropyl alcohol. Organisms were then separated from the remaining debris by flotation and hand picking, identified by experienced taxonomists, and enumerated. Quality assurance and control measures included randomly selecting 10% of the samples and reconstituting with the original debris and repeating the entire separation, identification, and enumeration process. The quality assurance plan used by BVA calls for all samples processed by a particular sorter to be redone if a random audit of 10% of that sorter's work shows any samples to differ by more than 5% from the original results. For this project, all discrepancies were within 5%. The taxonomists previously worked on several projects from the area and verified each other's identifications. When differences of opinion occurred or when significant doubt remained, outside specialists examined the specimens. BVA performed a review of all taxonomic identifications at the request of WES in 1997 and existing discrepancies were rectified. Identifications were made to the lowest practical identification level (LPIL) when not to the species level. Wet-weight biomass was determined to 0.01 mg after grouping the specimens by class within each station and after blotting to remove excess liquid.

Sediment samples were transferred to whirl-pack bags in the field. Between Spring 1994 and Fall 1995, all sediment samples were processed by Tierra Consulting, Inc. of Mobile, Alabama. The samples were thoroughly mixed in a stainless bowl and a 30-50 g subsample was washed with deionized water, dried and weighed. Coarse and fine fractions of the subsample were separated through a 4-phi sieve (0.0625 mm). Sediment texture of the coarse fraction was determined at 1-phi intervals using nested sieves (-2 to 4-phi, 4.00 to 0.0625 mm)

on a Ro-Tap apparatus. The weight of material collected on each sieve was recorded and used to compute mean and median grain size, and percentages by phi interval. No formal analyses of the fine fractions were done. After Fall 1995, all intertidal sediments were analyzed by Wetlands and Coastal Ecology Branch (U.S. Army Engineer Research Development Center). The only change in sample processing was the replacement of physical sieving of the sediments using the Ro-Tap apparatus by a wet-sieving technique. There is no discernable difference in the results produced by the two techniques. Nearshore sediment texture samples were analyzed in a similar manner with the exception that the fine fractions (silts and clays) were explicitly measured using a flotation technique (pipette method).

Water quality data were collected with a HydroLab® sensor during different components of the sampling program. Data reported here were compiled from values obtained during surf zone ichthyoplankton and beach seine sampling as well as nearshore infaunal sampling. Data include water temperature (°C), salinity (ppt), dissolved oxygen (mg/l), and pH. Turbidity (NTU) was also measured but will be reported in Chapters 5 and 7.

***Descriptive and Statistical Analyses:*** Numerical abundance, biomass, and taxa richness data were analyzed using Analysis of Variance (ANOVA) employing a two-way repeated measures design with sampling date as the repeated measure. Prior to testing, all data were converted to a per-m<sup>2</sup> basis, examined for normality and homogeneity of variance, and transformed ( $\log(x+1)$ ) where appropriate. Because multiple analyses were performed on data from each depth, it was necessary to adjust p values for multiple tests using the Bonferroni correction ( $p = 0.5/n$  where  $n =$  number of tests). Since a total of nine tests were analyzed (abundance, biomass, and taxa richness at MLW, MLW-1m, and Nearshore), a p value of 0.006 was necessary to indicate the presence of a statistically significant difference.

If a significant difference ( $p < 0.006$ ) was not detected for an effect, an *a posteriori* statistical power of the test was calculated. A power level of 75-80% was assumed to be necessary to indicate that no statistical difference was present. Where the interaction factor (e.g., Area x Date) was significant ( $p < 0.006$ ), main effects could not be directly interpreted (Zar, 1996). In these cases, linear contrasts were performed on appropriate pairs of means. Because the significance level for interpreting contrasts was also calculated using the Bonferroni correction for multiple tests it was necessary to restrict the number of tests performed to conserve power. Tests were restricted to data from sampling periods during and immediately after nourishment: September 1997, May 1998, September 1999 and May 2000. Since a total of four comparisons were made for each parameter, a p value of  $< 0.0125$  was required for statistical significance. In all but one case, reference area means were compared to that of the nourished area. For instance, the first comparison was between South Area MLW September 1997 and the data for MLW of the remaining two areas. The only exception to this procedure was for taxa richness data from MLW in May 2000. Since, there was no obvious difference between the Middle (Nourished) and one of the reference (North Area) means (Figure 2-11), but there was between these two values and that of the South Area (Reference), the Middle and North Area values were compared to that of the South Area. Where differences were detected

among main or interaction effects, the arithmetic means and standard errors were plotted. Taxonomic distributions of biomass and sediment texture data were examined graphically but not statistically analyzed.

Patterns in community species composition were examined using the multivariate ordination technique Nonmetric Multidimensional Scaling (NMDS). NMDS utilizes a ranking protocol to remove nonlinear trends in the same way ranking is used to remove non-normality in nonparametric ANOVA's. Since ordination techniques are sensitive to the impact of rare species, only taxa contributing between 0.5% (intertidal) and 1.0% (nearshore) of abundance from any area were incorporated into the analyses. The difference in selection criteria used for samples from the different depths was due to the low number of taxa in intertidal samples. Abundances were logarithmically transformed ( $\log x+1$ ) to reduce the influence of high dominance by one or more taxa. Prior to estimation of the contributions of individual taxa, the list of LPIL taxa was consolidated to remove duplicative listings. For instance, the mole crab was listed both as *Emerita talpoida* and *Emerita* (LPIL). These data were consolidated to form the single taxonomic listing *Emerita talpoida*. In this fashion, the intertidal taxa list was reduced to 80 taxa and the nearshore list to 225 taxa.

Temporal patterns in recovery of abundance, biomass, and taxa richness from monthly samples were examined by linear regression. Individual station data were expressed as a percentage of reference values for each date; after arcsin-square root transformation, data were regressed against time (days) since completion of nourishment. Outliers were identified by visual examination of residual plots and removed.

## Results

***Biannual Infaunal Abundance:*** A total of 132,518 individuals (average density =  $9,388/\text{m}^2$ ) were collected from the intertidal zone and an additional 298,999 animals ( $632/\text{m}^2$ ) from the nearshore subtidal (Table 2-1). The apparent anomaly between total numbers of animals collected and average abundance is due to differences in sample size. In the nearshore, a total of  $3 \text{ m}^2$  were collected during each sampling (3 areas x 10 stations x 1 sample x  $0.1 \text{ m}^2/\text{sample}$ ), whereas only  $0.79 \text{ m}^2$  was collected during each intertidal sampling (3 areas x 10 transects x 2 depths x 3 samples x  $0.0044 \text{ m}^2/\text{sample}$ )

The highest intertidal abundances occurred at MLW of the Middle Area (38,963 individuals, average =  $16,583/\text{m}^2$ ) and the lowest at MLW-1m of the South Area (16,360 individuals, average =  $7,097/\text{m}^2$ ). Nearshore abundances were highest at the North Area (137,503 individuals, average  $883/\text{m}^2$ ) and lowest at the South Area (35,669 individuals, average =  $250/\text{m}^2$ ).

Intertidal abundance was dominated by three taxa: Rhynchocoela (LPIL), the spionid polychaete *Scoelelepis squamata*, and Oligochaeta (LPIL). Rhynchocoels were the overall numerical dominant providing 51% of all animals collected, while the other two taxa supplied 35% and 8%, respectively (Table 2-2). Rhynchocoels comprised over 66% of all Middle Area

infauna but only 37% and 44%, respectively, of the North and South Areas. The most abundant taxa also occurred most frequently (Table 2-2). Percent occurrence (% of samples where a taxon was present) was very similar for all three dominants in all areas. Rhynchocoels were present in slightly more than 36% of all samples, while *S. squamata* and Oligochaeta (LPIL) occurred in 30% and 16%, respectively, of all samples. Oligochaetes were particularly important at the North Area where they made up 19% of the assemblage. The blue mussel, *Mytilus edulis*, was also present in 14% of all samples and comprised more than 1% of all animals collected. This taxon, a dominant on the hard substrate rocky groins along the beach, is dislodged by heavy wave action and accumulates on the beach, hence their presence in the intertidal samples. Since these animals are not indigenous to beach sands they were not included in subsequent analyses. The polychaete *Protodriloides* (LPIL) made up almost 1% of all animals collected and was found in 6% of the samples. It was most abundant at the North and South areas. The amphipod *Corophium tuberculatum*, the polychaete *Microphthalmus* (LPIL), *E. talpoida*, and *Microphthalmus* sp. G all contributed approximately 1% of total abundance in the North Area but not to overall intertidal abundance. The remaining taxa made up less than 1% of total abundance but were present in 1% or more of all samples overall or at one of the three area (Table 2-2). Of these taxa, only Spionidae (LPIL) was excluded from further analyses on the assumption that it was comprised mostly of damaged or small specimens of *S. squamata*.

Nearshore samples were overwhelmingly dominated (74%) by the wedge clam, *Donax variabilis* (Table 2-3). The second most abundant taxon, the polychaete *Magelona papillicornis*, comprised almost 6% of total abundance, while the clams *Tellina agilis* and *Spisula solidissima* contributed 3.6% and 2.1% respectively. Other taxa contributing approximately 1% or more to total abundance included the amphipods *Acanthohaustorius millsii* and *Psammonyx nobilis* and the polychaetes *Dispio uncinata* and *Asabellides oculata*. The taxon Ampharetidae (LPIL) was excluded from further analyses because it most likely represents small or damaged specimens of another dominant, *Ampharete americana*, or a mixture of species. *M. edulis* was excluded for reasons previously described. The remaining dominant taxa either contributed 1% or more to the abundance within a specific area or were present in 10% or more of the samples (Table 2-3). It is worth noting that most of the intertidal dominants are present in this list but in lower abundance or frequency of occurrence.

ANOVA of the abundance data revealed a significant ( $p < 0.006$ ) interaction effect (Area x Date) at all three depths and as a result none of the other effects are interpretable (Table 2-4). Linear contrasts of the selected interaction means indicated a significant difference ( $p < 0.0125$ ) between Middle Area MLW values and those of the reference sites in May 1998, however values at this site were higher than the reference areas and are not believed to represent an impact response (Figure 2-5). No significant ( $p > 0.00125$ ) linear contrast was found for MLW-1m abundances in May 2000, although abundance at this time appears to be lower at the nourished area than the reference areas (Figure 2-6). A significant ( $p < 0.00125$ ) linear contrast was found however, for Nearshore abundance in May 2000 (Figure 2-7). Abundance was lower in the nourished area than the reference areas. In the case of both the MLW-1m and Nearshore abundances, the May 2000 values are well within the range of

normal values for unimpacted conditions. For instance, the abundance at MLW-1m in May 2000 was nearly identical to values at the South Area in May 1995, the Middle Area in September 1995 and the North Area in September 1996 (Figure 2-6). Likewise, Middle Area Nearshore abundance in May 2000 was similar to previous May values at this area and actually higher than three of the previous May values for the North Area, the only completely unimpacted site in the study (Figure 2-7).

**Biannual Infaunal Biomass:** More than 12 kg of wet-weight biomass was found in the intertidal samples and nearly 33 kg in the nearshore samples (Table 2-1). MLW samples averaged over 150 g/m<sup>2</sup>, MLW-1m samples averaged 97 g/m<sup>2</sup> and nearshore samples averaged 99 g/m<sup>2</sup>. As with abundance, the apparent anomaly between total and average biomass/m<sup>2</sup> in intertidal and nearshore biomass comparisons is the result of differences in sample size. South Area MLW had the highest average biomass with over 220 g/m<sup>2</sup> and North Area Nearshore the lowest with 67 g/m<sup>2</sup>. Annelids composed 72% of total MLW biomass and 78% of MLW-1m biomass. Intertidal mollusc biomass was almost entirely composed of blue mussels (*Mytilus edulis*) washed off of the surrounding jetties rather than true members of the sandy beach infauna. When intertidal biomass is recalculated excluding the mussels, annelids comprised 83% and 78% of MLW and MLW-1m biomass, respectively and crustaceans made up an additional 16%-20%. Nearshore biomass was dominated by the molluscs *Tellina agilis* and *Spisula solidissima* making up 63.5% of the total (Table 2-1). Annelids contributed 10% and crustaceans made up 24% of nearshore biomass.

ANOVA for total wet-weight biomass detected a significant ( $p < 0.006$ ) interaction effect (Area x Date) for all three depths and as a result none of the other effects are interpretable (Table 2-4). Linear contrasts of during and immediately post-nourishment means indicated a significant difference ( $p < 0.0125$ ) between the nourished beach (South Area) and reference beaches in May 1998 for MLW (Figure 2-8) and Middle Area (nourished) and reference areas at all three depths in May 2000 (Figures 2-9 and 2-10). As in the abundance data, the difference between nourished and reference biomass in May 2000 represents an impact, however, the values are no lower than many of those from unimpacted conditions. Specifically, equivalent or lower biomass values were encountered at Middle Area MLW in September 1995 and again in May 1996, at the South Area in May 1994 and September 1996, and at the North Area in May of 1994, 1995, 1997 and 1998 (Figure 2-8). Biomass at MLW-1m of the nourished site was equal to or higher than that reported in the entire period between September 1995 and September 1996 (Figure 2-9). It was also equal to values reported from the South Area during September 1994, 1998, and 1999. In the nearshore, Middle Area biomass in 2000 was equal to or greater than that for the same area in May of all years except 1996 (Figure 2-10). It was also equal to or greater than that of South or North Area values in May 1996.

**Biannual Infaunal Taxa Richness:** Although diversity was not explicitly included as a test parameter in the original study design, this does not preclude its examination. Prior to examination, however, it is important to review the inherent limitations imposed by the study design. First and foremost, Jaramillo et al. (1995) have indicated that at a minimum of 4 m<sup>2</sup> is

required to account for 95% of all taxa present on most beaches. Since this level of effort exceeded the resources available, diversity comparisons will of necessity be relative. Likewise, since the number of taxa collected is a function of sample area, direct comparisons cannot be made between intertidal and nearshore sites due to the difference in gear types used in sampling. Taxa richness (number of taxa/sample) was selected as the test measure since diversity indices such as the Shannon-Weiner Index have been shown to be inadequate in analysis of beach nourishment projects (Nelson, 1985; Wilber and Stern, 1992). Such indices also should not be tested with parametric statistics due to an absence of information on their underlying mathematical distributions (Norris and Georges, 1993).

A grand total of the 80 LPIL taxa were collected at both intertidal depths and 225 in the nearshore zone (Table 2-1). Total numbers of taxa at intertidal depths tended to be higher at MLW-1m than MLW and decrease in value among areas along a gradient from south to north. This pattern was not present in the nearshore data where there was no pattern among areas.

As with abundance and biomass, ANOVA's for taxa richness among intertidal and nearshore data had significant ( $p < 0.006$ ) interaction effects (Table 2-4). Linear contrasts of MLW data indicated a difference among areas during the second fill operation, however the difference does not appear to be directly related to nourishment (Figure 2-11). Taxa richness was found to be higher at the South (reference) Area than either the Middle (nourished) or North (reference) Areas. Linear contrasts of both MLW-1m and Nearshore data indicated taxa richness were significantly ( $p < 0.0125$ ) lower at the Middle Area (nourished) than either reference area in May 2000 (Figures 2-12 and 2-13). Again, the low values were within the range of values encountered during non-impacted periods of time. At MLW-1m, values of taxa richness equal to that of at the nourished site in May 2000 occurred at the South Area in September 1998. In the nearshore, May 2000 values were equal to those occurring at the North Area in May 1995.

***Biannual Biomass Composition:*** Biomass composition (% of total biomass) varied among areas, depths, and sample dates. Annelids dominated biomass at both intertidal depths particularly in the spring (Figures 2-14 to 2-16). Crustacean biomass was generally second most important and was most prevalent in fall collections. Mollusc biomass was only dominant occasionally and then only in fall collections. Since most of this biomass was made up of *M. edulis* it's contribution can be considered to be minimal. Biomass composition appears to have been altered at the South Area during it's nourishment (September 1997) as evidenced by higher than normal proportions of molluscs at MLW and echinoderms at MLW-1m. There was no indication of a change in biomass composition after this time period or at anytime during the nourishment of the Middle Area in 1999 (Figures 2-17 to 2-19). Nearshore biomass was dominated by molluscs followed, in order of importance, by crustaceans and annelids (Figures 2-20 to 2-22). Molluscs were generally most dominant in fall collections while crustaceans and annelids were most dominant in the spring. There is no indication of a change in biomass composition associated with nourishment operations.

**Biannual Species Composition:** Patterns in species composition were explored using the ordination technique Nonmetric Multi-Dimensional Scaling (NMDS). Ordination of data from all three depths was successful (Figure 2-23) with stress, a measure of “goodness of fit”, low at a value of 0.125 (Clarke and Warwick, 1994). A stress value of 0.20 or greater indicates that the data cannot be interpreted (Clarke and Warwick, 1994). Data ordinated almost entirely based on depth; the intertidal samples ordinated high on Axis 1 while the nearshore samples ordinated high on Axis 2 (Figure 2-23). There were no apparent differences between intertidal depths or areas within depths. Taxa positively and significantly ( $r^2 \geq 0.4$ ) correlated with Axis 1 (i.e., those characteristic of the intertidal stations) included *Rhynchocoela* (LPIL), *Scolelepis squamata*, and *Oligochaeta* (LPIL) (Table 2-5). Taxa positively and significantly ( $r^2 \geq 0.4$ ) correlated with Axis 2 (i.e., those characteristic of the nearshore stations) included *Donax variabilis* and *Magelona papillicornis*.

In order to detect temporal changes in species composition, especially those potentially associated with nourishment, two further analyses were employed. First, NMDS was performed on data for each depth separately, however, none of the results could be interpreted (stress > 0.20). A second strategy was then employed, plotting the original NMDS results but excluding data points from periods of time when no nourishment was occurring. Specifically, only data from September 1997, May 1998, September 1999 and May 2000 were plotted (Figure 2-24). Examination of these data indicated several cases where a change in species composition was associated with nourishment. In September, South Area MLW samples differed from those of the Middle (Reference) Area. This pattern was repeated in September 1999 and May 2000 when samples and both Middle Area (Nourished) MLW and MLW-1m differed from the South Area (Reference). Examination of the individual species data for these time periods (Appendix Tables 2-2) suggests that the generally lower total abundance and taxa richness at these times are responsible for the observed differences rather than a change in the presence, absence, or relative abundance of any given taxon. In the nearshore samples, data from the South Area in September 1997 (during nourishment) and Middle Area in May 2000 (after nourishment) were different from the other samples collected during those time periods (Figure 2-24). Examination of individual species data (Appendix Table 2-3) indicates that abundance of *Donax variabilis* was most responsible for these differences. *Donax* was absent from the South Area samples in September 1997 while in May 2000 there were ten-times fewer clams in the Middle Area than either of the reference areas.

**Biannual Sediment Texture:** Sediment texture data were not contrasted statistically, however, average values are presented in this section. There were three distinct trends associated with the sediment composition results: depth, seasonal, and interannual (Figures 2-25 to 2-36). The most striking and consistent difference occurred between depths with finer sediments being associated with increased depth. The mean grain size (MGS) at MLW ranged between 0.27 mm and 1.78 mm with most values (28 out of 33) being greater than 0.40 mm (Figure 2-25). At MLW-1m MGS ranged from 0.18 mm to 1.16 mm with only 11 of 33 values greater than 0.40 mm (Figure 2-26), while in the Nearshore sediment MGS ranged between 0.05 mm to 0.22 mm (Figure 2-27). These differences are also reflected in the

proportions of individual grain size fractions. For example, in Spring 1996 more than 72% of MLW sediment at the South Area was composed of gravels and coarse to very coarse sands (Figure 2-28). Sediments at MLW-1m for this area contained only 60% of these materials while nearshore sediments contained less than 2% (Figure 2-31 and 2-34).

The next most pronounced difference among sediments was associated with the season of collection; sediments were generally coarser in the spring than in the fall. The coarseness of sediments in the spring reflects the general cycle of beach sediment erosion and deposition; erosion is most common during the winter and spring when strong storms are most frequent (Nordstrom, 1975). Large amounts of fine-grained sediments are removed from the beaches leaving relatively coarse sediments, while during the relatively mild summer months, fine sediments are deposited. This pattern is not absolute since the frequency and intensity of storm activity can change from year-to-year. During mild winters there may be no appreciable change in beach sediments.

The pattern of seasonally alternating coarseness in grain size was most pronounced in the MLW samples from the South Area (Figure 2-25). South Area MGS was far lower in spring than fall of 1994, 1996, and again in 1998. Similar results were found at South MLW-1 in 1996 and 1998 (Figure 2-27) and among North Area Nearshore sediments from 1996 to 1999 (Figure 2-28). These changes are also seen in the distribution of individual grain size fractions. For example, in 1994 the South Area MLW spring samples had a cumulative total of >50% very coarse sands and gravel, while in the fall, the cumulative total was approximately 10% (Figure 2-28). Similar patterns are present during most years for all depths and areas (Figures 2-29 to 2-36).

Interannual variation occurred in both the pattern of seasonal differences and in longer-term changes in sediment texture. The 1997 data are the most striking example of interannual differences in sediment composition. Sediments at both intertidal depths during May 1997 were finer than previous fall samples (September 1996) and as fine or finer than the succeeding September samples (Figures 2-25 to 2-26). A similar anomaly occurred in 1999 when all but Middle Area MLW-1m intertidal sediments were as fine or finer than the previous or succeeding fall. Presumably these results are due to relatively mild winters. The “fining” of the sediments does not appear to be related to beach nourishment activities since neither dredging or sand placement had begun in either case (May 1997 and May 1999). In addition, the effect was detected at all areas and not just the one designated for nourishment. Seasonal variation in sediment texture was present but less consistent in the Nearshore data. Spring sediments were as fine or finer than subsequent fall samples at the South Area in 1996 and 1998 and at the Middle Area in 1998 and 1999 (Figures 2-34 and 2-35). Spring samples were consistently coarser than fall samples in the North Area (Figure 2-36).

Sediment texture also varied between years. Intertidal sediments were particularly coarse in May of 1994, 1996, and 1998 with fine materials (silts, clays and fine and very fine sands) being more prevalent beginning in May 1997 (Figures 2-25 and 2-26). This pattern was

far more pronounced in the nearshore data where fine materials increased in proportion to other sediment fractions from May 1996 to September 1999. This pattern was reversed in May 2000 when silts, clays, and very fine sands declined in proportion to the other sediment fractions (Figures 2-34 to 2-36).

Likewise, a previously noted tendency for nearshore sediments to become progressively coarser along a South to North gradient (USACE, 1999) proved to be ephemeral. Between May 1995 and September 1997, fine sediment fractions, such as the fine and very fine sands, decreased in importance with increasing distance from Manasquan Inlet (Figures 2-34 to 2-36). The degree of difference between areas became progressively less over time, until they were roughly equal in May 1998. Afterwards, all three areas appear to follow similar trends and had nearly equal proportions of fine materials.

**Monthly Infaunal Abundance:** Data from the monthly samples were processed and analyzed in an identical fashion to that described for the biannual samples. Analysis of Variance for the 1997 data detected a significant ( $p < 0.017$ ) area by month interaction (Table 2-6). Linear contrasts of September, October, and November means found a significant ( $p < 0.017$ ) difference only in October, although nourished beach (South Area) abundances were lower than the reference area (Middle Area) in both October and November (Figure 2-37). Regression of monthly abundance (expressed as a percentage of reference values) produced a significant result;  $r^2$ -value = 0.633 and ANOVA  $p = 0.0059$  (Table 2-8). Estimated time for recovery of abundance based on this regression was 49 days.

Monthly abundance data for 1999-2000 yielded relatively high values during the spring and summer months, declining values during the winter, and lowest values in January (Figure 2-38). Analysis of Variance detected a significant ( $p < 0.017$ ) area by month interaction (Table 2-7). Of the three pairs of means tested (January, March and May), only the January data showed a significant difference ( $p < 0.017$ ). Abundance at the South Area (reference) was higher than that of the nourished Middle Area (Figure 2-38). Linear regression of the 1999-2000 abundance data produced a significant regression ( $p = 0.0026$ ), although the  $r^2$  value was only 0.24 (Table 2-8). The estimated time to recovery was 189 days or approximately 6.5 months.

**Monthly Infaunal Biomass:** Monthly biomass for 1997 monthly samples followed the same pattern as abundance with values declining with the onset of fall. ANOVA failed to detect any significant ( $p < 0.017$ ) differences (Table 2-6), however, power values were too low for all three effects (Area, Date, and Area x Date Interaction) to interpret the results. Highest biomass was present at the nourished area in September (Figure 2-39). This high value is attributable to the presence of two unidentified bivalves possibly washed onto the beach during the nourishment process (Appendix Table 2-5). The lowest biomass was found at the reference area in November while nourished area biomass was also low in both October and November. Regression of monthly biomass (expressed as a percentage of reference values) also produced a significant result;  $r^2$ -value = 0.547 and ANOVA  $p = 0.0145$  (Table 2-8). Estimated time for recovery of total biomass based on this regression was 38 days.

Monthly biomass for 1999-2000 also displayed a summer peak and winter low; lowest values occurred in January (Figure 2-40). ANOVA detected a significant ( $p < 0.017$ ) area by month interaction and of the three dates tested only January showed a significant linear contrast ( $p < 0.017$ ) (Table 2-7). Biomass was higher at the reference area than the nourished site at this time (Figure 2-40). Linear regression of the 1999-2000 biomass data produced a significant regression ( $p = 0.0016$ ) although the  $r^2$  value was only 0.24 (Table 2-8). Estimated time for recovery of total biomass based on the regression was 176 days or roughly 6 months.

**Monthly Infaunal Taxa Richness:** Taxa richness values for both sets of monthly data (1997 and 1999-2000) matched those of the abundance and biomass data with peak values in summer and lowest values in January (Figures 2-41 and 2-42). No significant differences ( $p > 0.017$ ) were found in the ANOVA of the 1997 data, however, statistical power was too low to interpret the results (Table 2-6). Regression of monthly taxa richness produced a significant ANOVA result ( $p = 0.0072$ ), with an  $r^2$  value of 0.616 (Table 2-8). Estimated time for recovery of taxa richness based on this regression was 23 days. ANOVA of the 1999-2000 data produced a significant interaction effect and subsequent linear contrasts found only the January comparison to be significant ( $p < 0.017$ ). As with abundance and biomass, taxa richness was lower at the nourished beach than the reference beach (Figure 2-39). Linear regression of the 1999-2000 data produced a significant regression ( $p = 0.0002$ ), but with a low  $r^2$  value ( $r^2 = 0.36$ ). Estimated time for recovery of total biomass based on this regression was 178 days or roughly 6 months (Table 2-8).

**Monthly Infaunal Biomass Composition:** In 1997, annelids dominated infaunal biomass in both areas during most collection periods. Crustaceans and molluscs dominated a few stations in August and again in November (Appendix Table 2-7). There was no obvious association between nourishment and the numbers of stations dominated by a particular taxonomic group or the extent to which a taxonomic group dominated biomass at a given station. For instance, in October 1997, when disturbance should have been at its peak, annelids comprised more than 95% of biomass at all stations in both areas. At the end of nourishment in November 1997 both areas had two stations dominated by annelids, two or three dominated by crustaceans, and one station with no biomass.

In 1999-2000, most stations in both areas were dominated by annelids between May and November 1999 (Appendix Table 2-8). In December 1999, stations at the reference beach (South Area) continued to be dominated by annelids while at the nourished beach (Middle Area), two stations had no biomass, one was dominated by annelids, one by molluscs, and two by miscellaneous taxa. The following January, four of the six nourished stations had no biomass, one was dominated by annelids and the remaining station was dominated by miscellaneous taxa. Annelids, crustaceans, and molluscs each dominated two reference area stations. During the remainder of the study (February 2000 to May 2000), annelids were dominant at the majority of stations in both areas. These results indicate a nourishment impact lasting from December 1999 to January 2000 with recovery complete by February 2000. While biomass composition results cannot be analyzed in the same fashion (linear regression) as abundance, biomass, and taxa richness data, a conservative estimate can be made by calculating

the time between completion of nourishment at the first station to be disturbed (September 27, 1999) and the February sampling (February 19, 2000) when recovery appears to have been complete. This calculation resulted in a recovery time of 145 days or approximately 4.8 months, a value close to the 176 day recovery estimate for total biomass.

**Monthly Infaunal Species Composition:** Because of the small number of taxa involved, NMDS could not be performed on the monthly infaunal samples. Instead, potential changes in species composition are assessed by directly examining taxa abundances (Appendix Tables 2-5 and 2-6). During 1997, rhynchocoels, *Scolelepis squamata*, and oligochaetes were the most abundant taxa at both sites and in all time periods (Appendix Table 2-5). There was no obvious change in species composition or the abundance of the dominant taxa related to nourishment.

In 1999-2000, rhynchocoels, *S. squamata*, *Protodriloides* (LPIL), oligochaetes and *Emerita talpoida* were the most abundant taxa (Appendix Table 2-8). Changes associated with nourishment included the absence of *Protodriloides* (LPIL) until May 2000 after nourishment and a reduction in the abundance of *S. squamata* relative to the reference area between November and December (Figure 2-43).

**Monthly Infaunal Sediment Composition:** During the 1997 nourishment, sediments at both the nourished beach (South Area) and reference beach (Middle Area) were characterized by mean grain sizes less than 0.5 mm throughout the study period (Figures 2-44 and 2-45). There was no obvious difference in sediment texture after nourishment. The same is true for the 1999-2000 nourishment (Figures 2-46 and 2-47). Mean grain sizes were generally less than 0.5 mm at both the nourished beach (Middle Area) and reference beach (South Area), although higher values were found at Station 1 (South Area) in May and July of 1999 and January of 2000. High values were also found at Station 2 in September 1999 and January 2000. There was no change in sediment texture associated with the 1999 nourishment.

**Water Quality Data:** Water quality parameters varied seasonally and among depths. As might be expected, water temperatures were higher in September than May of all years and generally 1-3 C° higher in the surfzone than in nearshore bottom waters (Figure 2-48). There were no consistent differences among areas and no obvious project-related effects during either dredging operation. Salinity varied among seasons and depths with lowest values occurring during May sampling periods (Figure 2-49). Salinity was usually 2-4 ppt lower in the surfzone than in nearshore bottom waters. Like the temperature results, there were no consistent differences among areas and no obvious changes due to dredging operations. Dissolved oxygen concentrations were similar in both the surfzone and nearshore bottom waters during spring sampling periods, but up to 4 mg/l lower in the nearshore waters during fall collections (Figure 2-50). This was particularly evident in September 1995 and again in September 1997. Comparatively low dissolved oxygen concentrations were also measured in bottom waters at the offshore borrow areas in September 1997 (See Chapter 8). There does not appear to be a relationship between the low values and nourishment since similar values were found in both nourished and unnourished areas and even lower values had been present when no nourishment

was going on (September 1995). Varying over a range of less than one unit, pH was generally higher in September than May sampling periods (Figure 2-51). There were no consistent differences among areas or depths, and no apparent changes related to either dredging operation.

## Discussion

Infaunal assemblages of high-energy sandy beaches are dominated by two different types of organisms: small interstitial forms including rhynchocoels, oligochaetes, and hesionid and protodile polychaetes, and large mobile forms such as the mole crab, *Emerita talpoida*, the wedge clam *Donax variabilis*, and the polychaete *Scolelepis squamata* (McLachlan and Jaramillo, 1995). Interstitial forms tend to dominate numerical abundance while the larger organisms dominate biomass. Infauna are distributed intertidally among three to four zones although it is difficult to precisely assign assemblages to individual zones because of the mobile nature of both the organisms and their environment. Species such as *Emerita talpoida* and *Donax variabilis* undergo diurnal and seasonal migrations (Bowman and Dolan, 1985; Eilers, 1995a,b), while the interstitial taxa are susceptible to being dislodged from the sediment by wave action or moved by bedload transport.

On the U.S. Atlantic coast the uppermost or supralittoral zone is dominated by air-breathing crustaceans such as talpid amphipods (also known as beach-hoppers) and ocypodid (ghost) crabs (McLachlan and Jaramillo, 1995). Between the drift line and midtide level is the littoral or swash zone, an area dominated by isopods, haustoriid amphipods and polychaetes such as *Scolelepis squamata*. Below midtide lies the sublittoral zone where *Emerita*, *Donax*, and a variety of haustoriid amphipods typify the benthic assemblage. On some beaches (macrotidal dissipative), littoral zone benthos may be further subdivided into assemblages associated with the resurgence and saturation zones described by Salvat (1964, 1967) (McLachlan, 1990 and Raffaelli, et al., 1991). Beyond the beach, both McLachlan et al. (1984) and Fleischack and de Freitas (1989) have identified benthic assemblages associated with the area of breaking waves (breaker zone) and a nearshore zone beyond the breakers. Assemblages of these zones are characterized by the decreasing representation of beach fauna and increasing importance of fauna characteristic of offshore waters.

The distribution of beach infauna is controlled by physical factors, particularly wave energy and tidal range (McLachlan, 1990), as manifested in beach morphology (Wright and Short, 1984; Short, 1991). Different combinations of wave energy and tidal range produce characteristic beach types which can be classified by their slope and the average height of incoming waves (Masselink and Short (1993). Beach slope and wave height, in turn, have been identified as the two factors associated the most with different beach assemblages (McLachlan 1990; McArdle and McLachlan 1991, 1992). Wave height is important because it is a measure of wave energy: the higher the wave energy, the more stressed and therefore less diverse and abundant the infaunal assemblage (McLachlan, 1983). Beach slope is important because beaches with steep slopes have a relatively small swash zone and species such as *Emerita* and *Donax* which “ride” the tides in the swash zone (Bowman and Dolan, 1985;

Ellers, 1995ab) may not have sufficient scope for feeding and thus be unable to establish large populations (Leber, 1982b). Lateral features of beach morphology also influence assemblage structure. McLachlan and Hesp (1984) have shown that meiofauna, macrofauna, and nekton were distributed in different manners along the cusps and horns of a cusped beach. Meiofauna were concentrated along the sides of the rip current, macrofauna in the cusp center, and nekton near the head of the rip currents.

Sediment texture, which is largely determined by the wave environment, can be a factor, although it is more difficult to detect unless the change in grain size is relatively large. Dexter (1969) and Leber (1982b) sampled beaches in the vicinity of Morehead City, North Carolina, but found relatively different assemblages. Dexter sampled a fine and medium sand beach dominated by haustoriid amphipods, while Leber found a medium sand beach to be dominated by *Emerita* and *Donax*. McLachlan (1996) has found that placement of coarse sands associated with mine tailings on a sandy beach in South Africa altered both beach morphology and the intertidal fauna. Species associated with the naturally occurring fine sands were replaced by larger, more robust fauna. In Uruguay, Defeo et al. (1997) have noted that the distribution of fine and coarse sands have substantial impacts on the distribution of species of the isopod genus *Exocirolana*.

Species composition in the study area was similar to that of other New Jersey and Atlantic coast beaches, although the relative dominance of individual taxa differed (Table 2-9). In an examination of sandy beach infauna in Maine, Larsen and Doggett (1990) found three different assemblages. Corresponding closely to temperature discontinuities, two assemblages were found north of Mount Desert; the northernmost was characterized by opheliid and paraoniid polychaetes and the other by oligochaetes and nepthyid and orbiniid polychaetes. The third and southernmost assemblage was dominated by *Scolelepis squamata* and the amphipods *Amphiporeia virginica* and *Haustorius canadensis*. This southernmost assemblage was the only one representing exposed (high-energy) beaches while the more northerly assemblages represent protected or low-energy beaches. Croker et al. (1975) and Croker (1977) described a similar distribution in southwestern Maine and New Hampshire. Amphipods dominated the more exposed sites, while polychaetes tended to be most abundant in areas with less direct exposure to wave action. In a study of sandy beach infauna near Avalon, New Jersey, McDermott (1983) reported *Scolelepis squamata* as the overall dominant followed by *Donax variabilis*, *Emerita talpoida*, *Amphiporeia virginica*, *Micrura leidy* (a rhyngocoel), and the haustoriid amphipod *Haustorius canadensis*. Five other haustoriids were also present at the site. Croker (1970) has reported the same assemblage from sandy beaches of Long Island, New York. As previously noted, Dexter (1969) has described a North Carolina beach community dominated (in order) by haustoriids, *Donax*, and *Scolelepis*, while Leber (1982b), working in the same general area, found *Emerita*, two species of *Donax*, and haustoriids to be most abundant. Diaz and DeAlteris (1982) reported similar distributions for a beach at Duck, North Carolina. In a study of a South Carolina beach Knott et al. (1983) found *Scolelepis* and haustoriid amphipods accounted for 63% of all animals present with *Donax* abundant in two of three sampling transects.

Intertidal infauna in the project reach of the present study were dominated by interstitial rhynchocoels, oligochaetes, hesionid and protodile polychaetes, and larger forms such as *Emerita talpoida*, *Scolelepis squamata* and *Nephtys buccera* (Table 2-2). The interstitial forms dominated abundance while the large forms dominated biomass. *Scolelepis* was unique in that it was dominant in both abundance and biomass. As previously mentioned, large numbers of the blue mussel, *Mytilus edulis*, were also present in the samples but were excluded from consideration since they were not true beach infauna. Nearshore fauna were dominated by the larger forms with *Donax variabilis* being especially abundant (Table 2-3). The tendency for *Donax* to be most abundant in the nearshore rather than intertidally may be a function of beach slope.

As noted in a previous report (USACE, 1999), the distribution of intertidal infauna in the study area was patchy with "hot-spots" of high abundance and/or biomass occurring unpredictably. High numerical abundances were generally associated with interstitial infauna such as rhynchocoels, enchytraeid oligochaetes, and archiannelids. These small animals live on and between sand grains and are routinely redistributed by wave action. Concentrations of suspended meiofauna can be considerable with up to a third of the total consisting of interstitial forms (e.g., Bell and Sherman 1980; Hagerman and Rieger 1981). Between 1994 and 1996 total assemblage numerical abundance was generally highest at MLW of the Middle Area in spring, however this pattern dissipated after 1996 (Figure 2-5). Peaks in abundance coincided with high rhynchocoel densities. There was no predictable difference in abundance among areas or seasons at the other depths. "Hot-spots" of biomass seemed to correspond to high densities of the mole crab *Emerita* and occasionally to abundance of *Scolelepis* in the intertidal and *Donax* in the nearshore. Both *Emerita* and *Donax* migrate up and down the swash zone during a tide so how many animals are encountered may vary with the time of sampling. Highest abundances of *Emerita* are associated with low wave energy and low tides (Bowman and Dolan 1985). Seasonally both abundance and biomass displayed peak values in the summer and fall and lowest values in mid-winter. This pattern has also been noted in Florida (Salomon and Naughton, 1984), South Carolina (Van Dolah et al., 1994; Jutte et al., 1999b), and North Carolina (Leber, 1982b; Reilly and Bellis, 1983).

While abundance values are low compared to those of McDermott (1983) at Avalon Beach, NJ, they are similar to densities reported from other Atlantic coast beaches (Table 2-9). It should also be noted that the present study encompasses a much larger stretch of beach and longer time period than that of McDermott (1983) and as a result incorporates a higher degree of spatial and temporal variability.

During the 1997 nourishment operation, no deleterious impacts to the intertidal assemblages were detected. The biannual (long-term) samples showed no indication of a difference in abundance between the nourished beach (South Area) and either of the reference areas (Figures 2-5 to 2-6). Biomass results were very similar with little predictable pattern in seasonal or geographic distribution from the biannual samples (Figures 2-8 to 2-9). Monthly data indicated very high biomass in September but low values afterwards (Figure 2-39). Diversity, as measured by taxa richness, did not appear to differ among areas or seasons and no

impact was detected associated with nourishment (Figures 2-11 to 2-12). Likewise, no dramatic differences could be detected in either biomass composition or species composition. Biomass was dominated either by annelids (mostly *Scolelepis*) or crustaceans (mostly *Emerita*) but without a distinct pattern of spatial or temporal distribution (Figures 2-14; 2-17; 2-20). The only changes in biomass composition associated with nourishment appeared to be the stranding of a few large bivalves on the nourished site immediately after nourishment. Likewise, there were no consistent differences in species composition among areas, depths or dates and no apparent change in assemblage structure following nourishment (Appendix Tables 2-2 and 2-5). The only discernable change in the nearshore assemblage was the temporary disappearance of *Donax variabilis* (Appendix Table 2-3).

During the 1999 nourishment operation, there were clear but short-lived impacts to abundance, biomass, and taxa richness at all three depths. Abundance and biomass, in particular, were still lower at the placement area than the reference sites as late as May 2000. Short-term (monthly) sampling at MLW indicated that the most severe impacts occurred in the period November 1999 to January 2000. Values began to reach reference site values only between February and April 2000 (Figures 2-38; 2-40; 2-42). The same pattern was evidenced in the abundance of *Scolelepis squamata* (Figure 2-43). Biomass composition did not change in respect to nourishment during this period. Changes in species composition in the intertidal zone were related to declines in abundance and taxa richness rather than an altered assemblage structure. In the nearshore there was no difference in species composition during nourishment, but afterwards abundances of *Donax* and *Asabellides oculata* were lower in the nourished area than the reference sites. Despite the continuing lower abundance and biomass in May 2000 and the occasional differences in the abundance of individual species, these results were within the range of values reported during unimpacted conditions. Based on the 6.5 month estimate for recovery time, approximately 80% of the nourishment area had already recovered and only the portion nourished after mid-November (Stations 21-23) was still impacted.

Results of both the long-term (biannual) and short-term (monthly) sampling are consistent with previous studies of beach nourishment impacts. Impacts tend to be most severe to small relatively immobile species, those unable to burrow through the overburden of new sand (e.g., Maurer et al. 1978). Impacts during the present study were most apparent to densities of rhynchozoels, oligochaetes, and the polychaete *Scolelepis squamata* (e.g., Figure 2-40). Fortunately, these taxa generally have high reproductive rates, wide dispersal capabilities, and can recover in short periods of time. Larger, more mobile taxa which can burrow through the new sediment or avoid the disturbance by migrating out of the area are generally less impacted by sediment deposition. Hayden and Dolan (1974) examined the impact of beach nourishment on the mole crab, *Emerita talpoida*, and found that while crab abundance declined in the immediate proximity of the nourishment there was no evidence of mortality; crab abundance recovered within a few weeks. The authors hypothesized that the crabs migrated out of the immediate area during the disturbance but rapidly colonized the site after the disturbance was over. Schoeman et al. (2000) have conducted manipulative experiments on beach infauna and reported that defaunated sediments were colonized within two weeks.

Literature reviews of beach nourishment impacts to beach infauna (Nelson, 1985, 1993 and Hackney et al., 1996) categorize impacts as short-term with recovery times ranging from 2 to 7 months. For instance, Saloman and Naughton (1984) reported infaunal recovery within 5-6 weeks of nourishment at Panama City, Florida. Studies by Van Dolah et al. (1994) and Jutte et al. (1999a,b) report recovery periods of 3 to 6 months for nourished beaches in South Carolina. The longest estimated recovery times for beach nourishment monitoring studies occurred when the silt/clay content of fill materials was higher than that of the natural beach. Reilly and Bellis (1983) reported that recovery took more than a year for some species on a North Carolina beach. Recovery was not complete until the silts and clays had been dissipated by wave action. Rakocinski et al. (1996) also found delayed recovery of infaunal assemblages due to the presence of substantial amounts of silts and clays in fill material used for a nourished beach and shallow inshore habitats in Perdido Key, Florida. Jaramillo et al. (1987) followed the natural recovery of a New England sandy beach after intense erosion and found that it took nearly three years for sand to accrete to pre-disturbance levels. During this time, polychaetes dominated the infauna; amphipod abundances did not reach pre-disturbance levels until a year later. Likewise, Peterson et al. (2000) found that the presence of finer than normal sediments and large amounts of shell hash in fill materials resulted in delayed recovery of large fauna such as *Emerita*, *Donax* and the ghost crab *Ocypode*. This study was further complicated by the fact that placement occurred during the late spring-early summer when filling could be expected to interfere with recruitment.

Estimates of recovery time in the present study, 2 months for the 1997 nourishment and 6.5 months for the 1999 operations, are precisely in the range of values found where there was a good match between fill materials and natural beach sediments (Table 2-10). The difference in recovery rates between the two nourishment operations seems most likely due to when placement was finished. The 1997 operation was completed by early October, whereas the 1999 nourishment was not completed until mid-December. Infaunal populations decline precipitously between November and January (e.g., Figure 2-40), suggesting that in 1997 there was enough time for colonization to be completed before the onset of the decline. In 1999, nourishment was not completed until a decline was well underway with the repercussion that there were insufficient animals available to fully colonize the disturbed sediments. The similarity between the results of this program and those of previous studies indicate that the findings will be applicable to subsequent renourishment operations at these sites and to similar projects in the New York-New Jersey area.

The principal conclusions from this portion of the study are as follows:

- 1) The intertidal infaunal assemblage was dominated by rynchocoels, the polychaetes *Scolelepis squamata*, *Protodriloides* (LPIL), and *Microphthalmus* spp., oligochaetes, the mole crab *Emerita talpoida*, as well as a number of haustoriid amphipods.

- 2) The nearshore infaunal assemblage included many of the same taxa, but was dominated by the wedge clam, *Donax variabilis*, the polychaete *Magelona papillicornis*, the clams *Spisula solidissima* and *Tellina agilis*, and the amphipods *Acanthohaustorius millsii* and *Psammonyx nobilis*, and the polychaete *Asabellides oculata*.
- 3) Infaunal assemblages of intertidal and nearshore beach environments were similar in species composition and abundance to those reported elsewhere on the Atlantic Coast. Abundance was somewhat lower than that reported for beaches in Southern New Jersey.
- 4) Intertidal abundances were highest in the summer and lowest in mid-winter.
- 5) Intertidal sediments varied between depths, seasons, and years. Mean grain size declined with depth and was generally highest in the spring.
- 6) Beach nourishment resulted in short-term declines in abundance, biomass, and taxa richness.
- 7) Recovery of intertidal assemblages was complete within 2-6.5 months of the conclusion of filling. Differences in the rate of recovery were most likely due to differences in when nourishment was complete. Sites where filling did not conclude until the low point in the seasonal cycle of infaunal abundance took the longest to recover.
- 8) Recovery rates are similar to those reported from other studies, particularly where the grain size of the fill material matched that of the beaches to be nourished.
- 9) There is no evidence of long-term impacts of beach nourishment operations on intertidal or nearshore infaunal assemblages.

## Literature Cited

- Bauer, B. O. and J. R. Allen. 1995. Beach steps: an evolutionary perspective. *Marine Geology* 123: 143-166.
- Bell, S. S. and K. M. Sherman. 1980. A field investigation of meiofaunal dispersal: tidal resuspension and implications. *Marine Ecology Progress Series* 3: 245-249.
- Bowman, M. L. and R. Dolan. 1985. The relationship of *Emerita talpoida* to beach characteristics. *Journal of Coastal Research* 1: 151-163.
- Cochran, W. 1963. *Sampling Techniques*, Second Edition. J. Wiley and Sons, Inc., New York, NY, 413 pp.
- Cohen, J. 1988. *Statistical Power Analysis for the Behavioral Sciences*, Second Edition. Lawrence Erlbaum Associates, Publishers, Hillsdale, NJ, 575pp.
- Clarke, K. R. and R. M. Warwick. 1994. Change in marine communities: an approach to statistical analysis and interpretation. Plymouth: Plymouth Marine Laboratory, 144pp.
- Croker, R. A. 1970. Intertidal sand macrofauna from Long Island, New York. *Chesapeake Science* 11: 134-137.
- Croker, R. A. 1977. Macro-infauna of northern New England marine sand: long-term intertidal community structure. pp. 439-450. In *Ecology of Marine Benthos*, B. C. Coull (Ed.). University of South Carolina Press, Columbia, SC.
- Croker, R. A., R. P. Hager and K. J. Scott. 1975. Macroinfauna of northern New England marine sand. II. Amphipod-dominated intertidal communities. *Canadian Journal of Zoology* 53: 42-51.
- Defeo, O., A. Brazeriro, A. de Alava and G. Riestra. 1997. Is sandy beach macrofauna only physically controlled? Role of substrate and competition in isopods. *Estuarine, Coastal and Shelf Science* 45: 453-462.
- Dexter, D. 1969. Structure of an intertidal sandy beach community in North Carolina. *Chesapeake Science* 10: 93-98
- Diaz, R. J. and J. T. DeAlteris. 1982. Long-term changes in beach fauna at Duck, North Carolina. U.S. Army Corps of Engineers Coastal Engineering Research Center Miscellaneous Report 82-12. November 1982. 48pp.
- Ellers, O. 1995a. Behavioral control of swash-riding in the clam, *Donax variabilis*. *Biological Bulletin* 189: 120-127.

- Ellers, O. 1995b. Discrimination among wave-generated sounds by a swash-riding clam. *Biological Bulletin* 189: 128-137.
- Fleischack, P. C. and A. J. de Freitas. 1989. Physical factors influencing the zonation of surf zone benthos. *Estuarine, Coastal and Shelf Science* 28: 517-530.
- Gorzelany, J. and W. Nelson. 1987. The effects of beach replenishment on the benthos of a sub-tropical Florida beach. *Marine Environmental Research* 21:75-94
- Hackney, C. T., M. H. Posey, S. W. Ross and A. R. Norris. 1996. A review and synthesis of data on surf zone fishes and invertebrates in the South Atlantic Bight and the potential impacts from beach nourishment. Prepared by the University of North Carolina at Wilmington and the North Carolina National Estuarine Research Reserve for the U.S. Army Corps of Engineers, Wilmington, NC.
- Hagerman, G. M. and R. M. Rieger. 1981. Dispersal of benthic meiofauna by wave and current action in Bogue Sound, North Carolina, USA. *P.S. Z.N. I Marine Ecology* 2: 245-270
- Hayden, B. and R. Dolan. 1974. Impact of beach nourishment on distribution of *Emerita talpoida*, the common mole crab. *J. Waterways, Harbors, and Coastal Engineering Division*, 100: 123-132.
- Jamarillo, E., R. A. Croker and E. B. Hatfield. 1987. Long-term structure, disturbance, and recolonization of macroinfauna in a New Hampshire sand beach. *Canadian Journal of Zoology* 65: 3024-3031.
- Jamarillo, E., A. McLachlan and J. Dugan. 1995. Total sample area and estimates of species richness in exposed sandy beaches. *Marine Ecology Progress Series* 119: 311-314.
- Jones, A. R., A. Murray, and R. E. Marsh. 1998. A method for sampling sandy beach amphipods that tidally migrate. *Marine and Freshwater Research* 49: 863-865.
- Jutte, P. C., R. F. Van Dolah and M. V. Levisen. 1999a. An environmental monitoring study of the Myrtle Beach Renourishment Project: Intertidal benthic community assessment of Phase I.- Cherry Grove to North Myrtle Beach. Final Report. Prepared by the Marine Resources Division, South Carolina Department of Natural Resources for U. S. Army Corps of Engineers, Charleston, SC.
- Jutte, P. C., R. F. Van Dolah and M. V. Levisen. 1999b. An environmental monitoring study of the Myrtle Beach Renourishment Project: Intertidal benthic community assessment of Phase II.- Myrtle Beach. Supplemental Report. Prepared by the Marine Resources Division, South Carolina Department of Natural Resources for U. S. Army Corps of Engineers, Charleston, SC.

Knott, D. M., D. R. Calder and R. F. VanDolah. 1983. Macrobenthos of sandy beach and nearshore environments at Murrells Inlet, South Carolina, U.S.A. *Estuarine, Coastal and Shelf Science* 16: 573-590

Larsen, P. F. and L. F. Doggett. 1990. Sand beach macrofauna of the Gulf of Maine with inference on the role of oceanic fronts in determining community composition. *Journal of Coastal Research* 6: 913-926.

Leber, K. M. 1982a. Bivalves (Tellinacea: Donacidae) on a North Carolina beach: contrasting population size structures and tidal migrations. *Marine Ecology Progress Series* 7: 297-301

Leber, K. M. 1982b. Seasonality of macroinvertebrates on a temperate, high wave energy sandy beach. *Bulletin of Marine Science* 32: 86-98

Masselink, G. and A. D. Short. 1993. The effect of tide range on beach morphodynamics and morphology: a conceptual model. *Journal of Coastal Research* 9: 785-800.

Matta, J. F. 1977. Beach fauna study of the CERC Field Research Facility, Duck, North Carolina. Miscellaneous Report 77-6. U. S. Army Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, Virginia, April 1977. 102p.

Maurer, D. L., Keck, R. T., Tinsman, J. C., Leathem, W. A., Wethe, C. A., Huntzinger, M., C. Lord and T. M. Church. 1978. Vertical migration of benthos in simulated dredged material overburdens. Volume 1: Marine benthos. Technical Report D-78-35. United States Army Corps of Engineers, Waterways Experiment Station, Vicksburg, MS.

McArdle, S. B. and A. McLachlan. 1991. Dynamics of the swash zone and effluent line on sandy beaches. *Marine Ecology Progress Series* 76: 91-99

McArdle, S. B. and A. McLachlan. 1992. Sand beach ecology: Swash features relevant to the macrofauna. *Journal of Coastal Research* 8: 398-407

McDermott, J. J. 1979. The role of a spionid polychaete in the food web of an intertidal community on an exposed sandy beach. *American Zoologist* 19: 865 (Abstract)

McDermott, J. J. 1983. Food web in the surf zone of an exposed sandy beach along the mid-Atlantic coast of the United States. pp. 529-538 in A. McLachlan and T. Erasmus (Eds.) *Sandy Beaches as Ecosystems*. Dr. W. Junk Publishers, Boston, MA.

McDermott, J. J. 1987. The distribution and food habits of *Nephtys buccera* Ehlers, 1868, (Polychaeta: Nephtyidae) in the surf zone of a sandy beach. *Proceedings of the Biological Society of Washington* 100: 21-27.

- McLachlan, A. 1983. Sandy beach ecology-a review. pp. 321-380 A. McLachlan and T. Erasmus (Eds.) *Sandy Beaches as Ecosystems*. Dr. W. Junk Publishers, Boston, MA.
- McLachlan, A. 1990. Dissipative beaches and macrofaunal communities on exposed intertidal sands. *Journal of Coastal Research* 6: 57-71
- McLachlan, A. 1996. Physical factors in benthic ecology: effects of changing sand particle size on beach fauna. *Marine Ecology Progress Series* 131: 205-217.
- McLachlan, A., A. C. Cockcroft, and D. E. Malan. 1984. Benthic faunal response of a high energy gradient. *Marine Ecology Progress Series* 16: 51-63.
- McLachlan, A. and P. Hesp. 1984. Faunal response to morphology and water circulation of a sandy beach with cusps. *Marine Ecology Progress Series* 19: 133-144.
- McLachlan, A. and E. Jaramillo. 1995. Zonation on sandy beaches. *Oceanography and Marine Biology: An Annual Review* 33: 305-335.
- Morrisey, D., L. Howitt, A. Underwood, and J. Stark. 1992. Spatial variation in soft-bottom benthos. *Marine Ecology Progress Series* 81:197-204.
- Nelson, W. G. 1985. Guidelines for beach restoration projects. Part 1. Biological. Florida Sea Grant Report 76. 66pp.
- Nelson, W. G. 1993. Beach restoration in the southeastern U.S.: environmental effects and biological monitoring. *Ocean and Coastal Management* 19:157-182
- Nordstrom, K. F. 1975. Beach response to cyclic and seasonal wave regimes at Sandy Hook, New Jersey. Technical Report 75-3. Marine Sciences Center, Rutgers- The State University of New Jersey, New Brunswick, NJ. pp. 57 & appendices.
- Norris, R. H. and A. Georges. 1993. Analysis and interpretation of benthic macroinvertebrate surveys. pp. 252-253 (Chapter 7) in D. M. Rosenberg and V. H. Resh (eds.) *Freshwater Biomonitoring and Benthic Macroinvertebrates*. Chapman and Hill, New York.
- Peterson, C. H., D. H. M. Hickerson and G. G. Johnson. 2000. Short-term consequences of nourishment and bulldozing on the dominant large invertebrates of a sandy beach. *Journal of Coastal Research* 16: 368-378.
- Raffaelli, D., I. Karakassis, and A. Galloway. 1991. Zonation schemes on sandy shores: a multivariate approach. *Journal of Experimental Marine Biology and Ecology* 148: 241-253.
- Rakocinski, C. F., Heard, R. W., LeCroy, S. E., McLelland, J. A. and T. Simons. 1996. Responses by macrobenthic assemblages to extensive beach restoration at Peridido Key, Florida, U.S.A. *Journal of Coastal Research* 12: 326-353.

Reilly, F. J. and V. J. Bellis. 1983. The ecological impact of beach nourishment with dredged materials on the intertidal zone at Bogue Banks, North Carolina. Miscellaneous Report 83-3. U.S. Army Engineer Coastal Engineering Research Center, Fort Belvoir, VA.

Saila, S., R. Pikanowski, and D. Vaughan. 1976. Optimum allocation strategies for sampling benthos in the New York Bight. *Estuarine, Coastal, and Marine Science* 4:119-128.

Saloman, C. H. and S. P. Naughton. 1984. Beach restoration with offshore dredged sand: effects on nearshore macrofauna. NOAA Technical Report NMFS-SEFC-133.

Salvat, B. 1964. Les conditions hydrodynamiques interstitielles meubles intertidaux et la repartition verticale del la fauna endogenee. *Academie des Sciences (Paris), Comptes Rendus* 259: 1576-1579.

Salvat, B. 1967. Les macrofauna carcinologique des sediments meubles intertidaux (tanaidaces, isopodes, et amphipodes): ethologie, bionomie et cycle biologique. *Memoires de la Museum National d'Historie Naturelle Series A* 45: 1-275.

Sameoto, D. C. 1969. Comparative ecology, life histories and behavior of intertidal and sand-burrowing amphipods (Crustacea: Haustoriidae) at Cape Cod. *Journal of the Fisheries Research Board of Canada* 26: 361-388.

Schoeman, D. S., A. McLachlan, and J. E. Dugan. 2000. Lessons from a disturbance experiment in the intertidal zone of an exposed sandy beach. *Estuarine, Coastal, and Shelf Science* 50: 869-884.

Short, A. D. 1991. Macro-meso tidal beach morphodynamics- an overview. *Journal of Coastal Research* 7: 417-436.

Underwood, A. 1992. Beyond BACI: the detection of environmental impacts on populations in the real, but variable, world. *Journal of Experimental Marine Biology and Ecology* 161:145-178.

USACE, 1999. The New York District's Biological Monitoring Program for the Atlantic Coast of New Jersey, Asbury Park to Manasquan Section Beach Erosion Control Project. Phases II-III. During Construction and 1<sup>st</sup> Year Post-construction Studies. U.S. Army Engineer District, New York.

Van Dolah, R. F., R. M. Martore, A. E. Lynch, P. A. Wendt, M V. Levisen, D. J. Whitaker, and W. D. Anderson, 1994. Final Report. Environmental Evaluation oof the Folly Beach Nourishment Project. Prepared by the Marine Resources Division, South Carolina Department of Natural Resources for U. S. Army Corps of Engineers, Charleston, SC. 101 p.p. & appendices.

Wilber, P. and M. Stern. 1992. A re-examination of infaunal studies that accompany beach renourishment studies. pp. 242-257 in *New Directions in Beach Management. Proceedings of the 5th Annual National Conference on Beach Preservation Technology*. The Florida Shore and Beach Preservation Association.

Wright, L. D. and A. D. Short. 1984. Morphodynamic variability of surf zones and beaches: a synthesis. *Marine Geology* 59: 93-118.

Zar, J. H., 1996. *Biostatistical Analysis*. 3d Ed. Prentice-Hall. 662p.